

Article

Cuticular Chemistry of the Queensland Fruit Fly *Bactrocera tryoni* (Froggatt)

Soo J. Park ^{1,2,*} , Gunjan Pandey ^{1,3} , Cynthia Castro-Vargas ^{1,3}, John G. Oakeshott ^{1,3} , Phillip W. Taylor ^{1,2}  and Vivian Mendez ^{1,2} 

¹ Applied BioSciences, Macquarie University, North Ryde, NSW 2109, Australia; Gunjan.Pandey@csiro.au (G.P.); Cynthia.Castro-Vargas@csiro.au (C.C.-V.); John.Oakeshott@csiro.au (J.G.O.); phil.taylor@mq.edu.au (P.W.T.); vivian.mendez@mq.edu.au (V.M.)

² Australian Research Council Centre for Fruit Fly Biosecurity Innovation, Macquarie University, North Ryde, NSW 2109, Australia

³ Commonwealth Scientific and Industrial Research Organisation Land and Water, Black Mountain, Acton, ACT 2601, Australia

* Correspondence: soojean.park@mq.edu.au; Tel.: +61-413-616-107

Received: 8 August 2020; Accepted: 10 September 2020; Published: 12 September 2020



Abstract: The cuticular layer of the insect exoskeleton contains diverse compounds that serve important biological functions, including the maintenance of homeostasis by protecting against water loss, protection from injury, pathogens and insecticides, and communication. *Bactrocera tryoni* (Froggatt) is the most destructive pest of fruit production in Australia, yet there are no published accounts of this species' cuticular chemistry. We here provide a comprehensive description of *B. tryoni* cuticular chemistry. We used gas chromatography-mass spectrometry to identify and characterize compounds in hexane extracts of *B. tryoni* adults reared from larvae in naturally infested fruits. The compounds found included spiroacetals, aliphatic amides, saturated/unsaturated and methyl branched C₁₂ to C₂₀ chain esters and C₂₉ to C₃₃ normal and methyl-branched alkanes. The spiroacetals and esters were found to be specific to mature females, while the amides were found in both sexes. Normal and methyl-branched alkanes were qualitatively the same in all age and sex groups but some of the alkanes differed in amounts (as estimated from internal standard-normalized peak areas) between mature males and females, as well as between mature and immature flies. This study provides essential foundations for studies investigating the functions of cuticular chemistry in this economically important species.

Keywords: cuticular hydrocarbons; cuticle; chemical communication; GC-MS; methyl branched alkanes; chemical ecology; volatiles

1. Introduction

The cuticular layer of the insect exoskeleton contains a range of mostly aliphatic compounds, including normal and branched alkanes, alkenes, saturated and unsaturated esters, alcohols, saturated and unsaturated fatty acids, ketones, and aldehydes [1]. Cuticular hydrocarbons usually contain 20 to 50 carbons, and compounds with other functional groups vary from 12 to 54 carbons [2–4]. A primary function of cuticular hydrocarbons is to protect against desiccation [5,6], injury, and infection [7–12]. Cuticular compounds, including hydrocarbons, are also commonly important for chemical communication [2,5,6,13], including species recognition [14–17], mimicry [18], and as pheromones [19] in diverse insect taxa. For example, some cuticular hydrocarbons serve as sex pheromones in house fly [20], the circumboreal fly [21], moths [22], bees [23] and the cowpea weevil [24]. Cuticular hydrocarbons also serve as aggregation pheromones in some insects, including

Drosophila [25], termites [26] and cockroaches [27]. Sexual selection has been a driving force for the evolution of sexual dimorphism in animals [28], and many insect taxa exhibit sexual dimorphism in cuticular chemistry [29]. For example, sexually dimorphic cuticular hydrocarbons have been found in *Drosophila* [30–32] and have been implicated in female attractiveness and male mating success [21].

Tephritid fruit flies are amongst the world's most economically damaging insect pests [33]. Some aspects of tephritid fruit fly semiochemistry have received significant attention, particularly the pheromones [34,35] they use to attract mates and for aggregation and the particular compounds found in fruit, food and certain flowers to which they are attracted [36–47]. Some work has also been performed on fruit fly cuticular chemistry because their cuticular chemical profiles tend to be highly species-specific [48,49] and have been used to resolve species, cryptic species and geographic variation in larvae [50–52] and adults [53–58]. Beyond their use as chemotaxonomic tools, however, relatively little work has been performed on tephritid cuticular compounds. In an important recent exception, allyl-2,6-dimethoxyphenol has been proposed as a short-range male attractant in *Bactrocera dorsalis* [59]. Most cuticular compounds are aliphatic, so this case is also notable for its involvement of an aromatic compound.

In Australia and in some Pacific Islands, the Queensland fruit fly, *Bactrocera tryoni* (Froggatt), is an economically important pest of horticultural crops [60–62]. This species causes significant economic loss by damaging crops [63] and by limiting market access [64]. While rectal gland and volatile emission chemistry of *B. tryoni* has been documented [65–70], the cuticular chemistry of this species has not. Given what is known for other insects, the composition of its cuticular chemical profile is likely to be relevant to homeostasis, protection from pathogens, injury and insecticides [71], and chemical communication. Understanding elements of cuticular chemistry related to homeostasis may help to understand abiotic factors mediating bioclimatic potential of *B. tryoni* [72] and effects of domestication, sex and age on desiccation resistance [73,74], and may also be important for understanding environmental and sexual competence of sterile *B. tryoni* released in sterile insect technique (SIT) programs to control pest populations [75–77]. To address this knowledge gap, and to provide foundations for subsequent functional studies, the present study reports a qualitative description of *B. tryoni* cuticular chemistry and identifies qualitative and quantitative variation (the latter estimated from internal standard-normalized peak areas) related to maturity and sex. *Bactrocera tryoni* specimens were obtained as larvae in infested fruits and cuticle extracts of emerged adults were analyzed by gas chromatography-mass spectrometry (GC-MS).

2. Results

Cuticular Chemistry and Statistical Analysis

The identified compounds are all aliphatic; no trace of aromatic compounds was found. Typical chromatograms of both immature and mature female and male *B. tryoni* are shown in Figure 1. The chromatogram sections of shorter (A) and longer retention time compounds (B) of *B. tryoni* adults are shown in Figure 2, where a typical chromatogram of a female is presented because the chromatogram includes all the compounds that are also found in immature and mature females and males. A range of non-alkanes, including spiroacetals, amides and esters and an assortment of C₂₉ to C₃₃ methyl-branched alkanes represent the cuticular chemistry of wild *B. tryoni* adults.

The identities of 22 of the 32 non-alkane compounds found were confirmed with authentic standards and the other ten were tentatively identified by comparison to fragment patterns in NIST libraries (Table 1). Two of the compounds were 6,6-membered ring spiroacetals (compounds A2 and A5 in Table 1 and Figure 2), four were aliphatic amides (compounds A1, A3, A4 and A6 in Table 1 and Figure 2), and the remaining 26 were all esters of saturated/unsaturated and methyl branched saturated fatty acids (compounds A7–A32 in Table 1 and Figure 2). The spiroacetals and saturated/unsaturated and branched saturated esters were found to be specific to mature females, while the amides were found in mature flies of both sexes (Table 1). In mature females, ethyl esters of saturated or unsaturated

C₁₂, C₁₄, C₁₆ and C₁₈ are the most abundant, while methyl and propyl esters, and branched saturated fatty acid esters are minor and trace, respectively. Methyl positions in the branched fatty acid esters are ambiguous, because trace amounts of the compounds made it difficult for further analyses. The amount of *N*-(3-methylbutyl)isobutyramide (**A6**) was about 2.5 times larger in mature male than in mature females ($p < 0.05$, *t*-test), but the amount of *N*-(3-methylbutyl)propanamide (**A4**) was about 3.2 times larger in mature females than in mature males ($p < 0.05$, *t*-test). The differences in the amounts of the other amides, *N*-(3-methylbutyl)acetamide (**A1**) and *N*-(2-methylbutyl)propanamide (**A3**) were not significant ($p > 0.05$ for all, *t*-test). The total amount of the amides was 2.6 times larger in mature females than in mature males ($p < 0.05$, *t*-test). The results are illustrated in Figure 3.




Figure 1. Typical chromatograms of hexane extracts of immature and mature female and male *B. tryoni*. (A) immature female; (B) immature male; (C) mature female; (D) mature male.




Figure 2. Representative chromatogram sections. (A) Non-alkanes in a chromatogram of a mature female *B. tryoni*, which includes all compounds found in mature and immature males and females; (B) Hydrocarbons (alkanes) section of chromatogram from a mature female *B. tryoni*. Note that compounds in B are qualitatively identical in immature and mature females and males.

Table 1. The compounds identified in hexane washes of *B. tryoni* that eluted early in chromatograms (section A in Figure 2).

No	Identity	MM	KI	Ref.KI (Ref)	Characteristic/Diagnostic EI Ions
A1 *	<i>N</i> -(3-Methylbutyl)acetamide	129.12	1131	1137 [78]	129 (M ⁺), 114, 86, 73, 60 (CH ₃ COHNH ₂ ⁺)
A2	2,8-Dimethyl-1,7-dioxaspiro[5,5]undecane	184.15	1140	1147 [78]	184 (M ⁺), 140, 115/112 (M-C ₅ H ₈ /C ₅ H ₈ O), 97, 69
A3 *	<i>N</i> -(2-Methylbutyl)propanamide	143.13	1198		143 (M ⁺), 114, 86, 74, 57
A4 *	<i>N</i> -(3-Methylbutyl)propanamide	143.13	1204		143 (M ⁺), 128 (M-CH ₃), 114 (M-C ₂ H ₅), 100, 87, 74, 57
A5 #	2-Ethyl-8-methyl-1,7-dioxaspiro[5,5]undecane	198.16	1230	1237 [78]	198 (M ⁺), 169, 129/126 (M-C ₅ H ₈ /C ₅ H ₈ O), 115/112 (C ₆ H ₁₀ /C ₆ H ₁₀ O), 97, 83, 69, 55
A6 *	<i>N</i> -(3-Methylbutyl)isobutyramide	157.15	1233		157 (M ⁺), 142, 101, 71, 57
A7	Ethyl dodecanoate (ethyl laurate)	228.38	1591	1593 [79]	228 (M ⁺), 183, 157, 115, 101, 88, 73, 70, 60 (CH ₃ CO=OH ⁺)
A8 #	Ethyl 6-methyldodecanoate	242.22	1662		242 (M ⁺), 213, 199, 185, 157, 143, 101, 88, 83, 70, 55
A9	Propyl dodecanoate	242.22	1680	1685 [80]	242 (M ⁺), 201, 183, 157, 143, 115, 102, 61 (C ₃ H ₇ OH ₂ ⁺ , base peak)
A10	Ethyl tridecanoate	242.22	1691	1695 [81]	242 (M ⁺), 199, 197, 157, 101, 88
A11	Methyl tetradecanoate	242.22	1722	1724 [82]	242 (M ⁺), 157, 143, 101, 87, 74
A12	Ethyl (<i>E</i>)-9-tetradecenoate (ethyl myristolaidate)	254.22	1769		254 (M ⁺), 208/209 (loss of EtOH/EtO), 166, 124, 88, 55
A13	Ethyl (<i>Z</i>)-9-tetradecenoate (ethyl myristoleate)	254.22	1778		254 (M ⁺), 208/209 (loss of EtOH/EtO), 166, 124, 88, 55
A14	Ethyl tetradecanoate (ethyl myristate)	256.43	1790	1793 [80]	256 (M ⁺), 213, 157, 101, 88
A15 #	Ethyl 4-methyltetradecanoate	270.26	1836		270 (M ⁺), 213, 101 (M-C ₁₂ H ₂₅ , base peak), 88
A16 #	Ethyl 12-methyltetradecanoate	270.26	1862		270 (M ⁺), 227, 213, 157, 101, 88
A17 #	Propyl tetradecanoate	270.26	1887	1893 [83]	270 (M ⁺), 229, 211, 172, 129, 102, 61 (C ₃ H ₇ OH ₂ ⁺ , base peak)
A18	Ethyl pentadecanoate	270.26	1890	1897 [84]	270 (M ⁺), 227, 199, 157, 101, 88
A19	Methyl (<i>Z</i>)-9-hexadecenoate	268.44	1902	1909 [78]	268 (M ⁺), 236/237 (loss of MeOH/MeO), 194, 152, 96, 74, 55
A20	Methyl hexadecanoate	270.26	1923	1927 [85]	270 (M ⁺), 227, 143, 87, 74
A21	Ethyl (<i>E</i>)-9-hexadecenoate (ethyl palmitelaidate)	282.26	1965		282 (M ⁺), 236/237 (loss of EtOH/EtO), 194, 152, 96, 88, 69, 55
A22	Ethyl (<i>Z</i>)-9-hexadecenoate (ethyl plamitoleate)	282.26	1970	1975 [86]	282 (M ⁺), 236/237 (loss of EtOH/EtO), 194, 152, 96, 88, 69, 55
A23	Ethyl hexadecanoate (ethyl palmitate)	284.27	1990	1993 [80]	284 (M ⁺), 241, 157, 101, 88
A24 #	Ethyl 15-methylhexadecanoate	298.29	2029		298 (M ⁺), 255, 157 (M-C ₁₀ H ₂₁), 101, 88
A25 #	Ethyl 4-methylhexadecanoate	298.29	2035		298 (M ⁺), 241 (M-C ₄ H ₉), 101 (base peak), 88
A26 #	Ethyl 14-methylhexadecanoate	298.29	2062		298 (M ⁺), 269, 255, 241, 199, 157, 101, 88
A27 #	Propyl 9-hexadecenoate	296.27	2067		296 (M ⁺) 281, 237, 194 (M-C ₃ H ₇ COOCH ₂)
A28	Ethyl (<i>Z,Z</i>)-octadeca-9,12-dienoate (ethyl linoleate)	308.27	2158	2155 [87]	308 (M ⁺), 262/263 (loss of EtOH/EtO), 178, 135, 95, 81
A29	Ethyl (<i>Z</i>)-9-octadecenoate (ethyl oleate)	310.29	2168	2168 [88]	310 (M ⁺), 264/265 (loss of EtOH/EtO), 222, 180, 97, 55
A30	Ethyl (<i>E</i>)-9-octadecenoate (ethyl elaidate)	310.29	2171	2174 [89]	310 (M ⁺), 264/265 (loss of EtOH/EtO), 222, 180, 97, 55
A31	Ethyl octadecanoate	312.54	2190	2191 [90]	312 (M ⁺), 269, 157, 101, 88
A32 #	Ethyl 11-eicosenoate	338.57	2366		338 (M ⁺), 292/293 (M-EtOH/EtO), 250, 208, 97, 55

MM: molecular mass, KI: Kovats' retention index, Ref.KI (Ref): reference KI if available for a similar column type, active phase and temperature conditions, with references in parentheses; * indicates that the compound is present in both sexes of mature adults, all other compounds being mature female specific; # indicates the compound was only tentatively identified.




Figure 3. Internal standard-normalized peak areas of *N*-(3-methylbutyl)acetamide (A1), *N*-(2-methylbutyl)propanamide (A3), *N*-(3-methylbutyl)propenamide (A4) and *N*-(3-methylbutyl)isobutyramide (A6) in section A in Figure 2 in female and male *B. tryoni*. Standardized peak areas were obtained by dividing the peak area of a compound by the peak area of the *n*-hexadecane internal standard. Error bars represent standard errors. The results of *t*-test comparisons between the sexes are shown (ns is not significant; * $p < 0.05$, ** $p < 0.01$).

The 34 tentatively identified hydrocarbons, all methyl-branched alkanes with C₂₉ to C₃₃ carbon backbones, are summarized in Table 2. Unsaturated hydrocarbons were not detected. Most are mono- or dimethylalkanes, with only a few trimethylalkanes found. Monomethyl branches appeared exclusively at odd carbon positions in odd carbon alkanes and at even carbon positions in even carbon alkanes. Comparisons in the amounts of individual hydrocarbons between sexes are illustrated in Figure 4. The most abundant alkanes were mono- and dimethylhentriacontane (C₃₁) isomers (compounds B18 to B24 in Figures 2B and 4), with the 11-, 13- and 15-methylhentriacontanes (B18) appearing at the highest intensity in chromatograms. Although dimethyl branches were separated by 1, 3, 5, 7, 9, 11, or 13 methylene groups, a majority of dimethyl branches were separated by 3 and 5 methylenes. Trimethyl branches were separated by [3,3], [3,5] or [3,11] methylenes.

There were significant effects of sexual maturity, sex and the interaction between these two variables on the amounts of the normal and methyl branched alkanes (sexual maturity $F_{1,2276} = 81.90$, $p < 0.0001$; $F_{1,2276} = 10.56$, $p < 0.005$; sexual maturity * sex $F_{1,2276} = 4.10$, $p < 0.05$, respectively). The amounts of alkanes in mature flies were 2.3 (± 1.3) times larger than those in immature flies on average. The amounts of compounds B1, B3, B4, B5, B6, B7, B9, B10, B11, B12, B13, B18, B19, B20, B21, B22, B22, and B24 were higher in mature females than in mature males ($p < 0.05$, *t*-test). The amounts of these compounds were 1.7 (± 0.5) times greater in mature females than in mature males on average. There was no significant difference in the amount of hydrocarbons between immature females and immature males ($p > 0.05$, *t*-test).

Table 2. The tentatively identified cuticular hydrocarbons found in *n*-hexane washes of *B. tryoni* (section B in Figure 1).

No	Identity	MM	KI	Ref.KI (Ref)	Characteristic/Diagnostic EI Ions
B1	11-; 13-; 15-MeC ₂₉	422.82	2929	2932 [91]	280/281,168/169; 252/253, 196/197; 224/225(s)
B2	7-MeC ₂₉	422.82	2946	2940 [51]	336/337, 112/113
B3	5-MeC ₂₉	422.82	2952	2949 [91]	364/365, 84/85
B4	9,13-DiMeC ₂₉	436.50	2966	2963 [91]	322/323, 140/141, 252/253, 210/211
B5	7,11-DiMeC ₂₉	436.50	2970		350/351, 112/113, 280/281, 182/183
B6	3-MeC ₂₉	422.82	2976	2973 [91]	392/393, 56/57
B7	5,11-DiMeC ₂₉ ; 5,13-DiMeC ₂₉	436.50	2986	2983 [91]	378/379, 84/85,280/281, 182/183; 378/379, 84/85, 280/281, 210/211
B8	4, <i>x</i> , 22-TriMeC ₂₉ (<i>x</i> = 14 or 16)	450.52	3009		392/393, 84/85, 252/253, 224/225, 336/337, 126/127
B9	12-Me; 14-MeC ₃₀	436.50	3025	3031 [92]	280/281,182/183; 252/253, 210/211
B10	8-MeC ₃₀	436.50	3034	3040 [93]	336/337, 126/127
B11	6-MeC ₃₀	436.50	3041	3045 [93]	364/365, 98/99
B12	4-MeC ₃₀	436.50	3055	3065 [93]	392/393, 70/71
B13	8,12-DiMeC ₃₀ ; 8,14-DiMeC ₃₀	450.52	3061	3064 [94]	350/351, 126/127, 280/281, 196/197; 350/351, 126/127, 253/252, 225.224
B14	6,14-DiMeC ₃₀ ; 6,12-DiMeC ₃₀	450.50	3071		378/379, 98/99, 252/253, 224/225; 378/379, 98/99, 280/281, 196/197
B15	4,12-DiMeC ₃₀ ; 4,14-DiMeC ₃₀ ; 4,20-DiMeC ₃₀	450.52	3088	3098 [94]	406/407, 70/71, 280/281, 196/197; 406/407, 70/71, 225/224, 253/252; 406/407, 70/71, 309/308, 169/168
B16	<i>n</i> -C ₃₁	436.50	3100		436
B17	4,8,12-TriMeC ₃₀ ; 4,8,14-TriMeC ₃₀ ; 4,8,20-TriMeC ₃₀	464.53	3115		70/71, 420/421, 350/351, 140/141, 280/291, 210/211; 70/71, 420/421, 350/351, 140/141, 252/253, 238/239; 70/71, 420/421, 350/351, 140/141, 322/323, 168/169
B18	11-; 13-; 15-MeC ₃₁	450.52	3129	3130 [92]	308/309, 168/169; 280/281, 196/197; 250/251, 224/225
B19	7-MeC ₃₁ ; 9-MeC ₃₁	450.52	3137	3140 [93]	364/365, 112/113; 336/337, 141/140
B20	5-MeC ₃₁	450.52	3147	3150 [93]	392/393, 84/85
B21	11,15-DiMeC ₃₁	464.53	3153	3155 [93]	322/323, 168/169, 252/253, 238/239
B22	9,13-DiMeC ₃₁ ; 9,15-DiMeC ₃₁ ; 11,13-DiMeC ₃₁ ; 13,15-DiMeC ₃₁	464.53	3157	3159 [92]	350/351, 140/141, 280/281, 210/211; 350/351, 140/141, 252/253, 238/239; 322/323, 168/169, 280/281, 210/211; 294/295, 196/197, 252/253, 238/239
B23	7,13-DiMeC ₃₁ ;7,15-DiMeC ₃₁	464.53	3164		378/379, 112/113, 280/281, 210/211; 378/379, 112/113, 252/253, 238/239
B24	3-MeC ₃₁	450.52	3170	3172 [91]	420/421, 56/57
B25	5,11-DiMeC ₃₁ ; 5,13-DiMeC ₃₁	464.53	3178	3180 [95]	406/407, 84/85, 182/183, 308/309; 406/407, 84/85, 280/281, 210/211
B26	12-; 14-; 16-MeC ₃₂	464.53	3223	3225 [57]	308/309, 182/183; 280/281, 210/211; 252/253, 238/239
B27	8-; 10-MeC ₃₂	464.53	3225	3225 [57]	364/365, 126/127; 336/337, 154/155
B28	10,14-DiMeC ₃₂	478.55	3233	3254 [94]	350/351, 154/155, 280/281, 224/225

Table 2. Cont.

No	Identity	MM	KI	Ref.KI (Ref)	Characteristic/Diagnostic EI Ions
B29	8,12-DiMeC ₃₂ ; 8,14-DiMeC ₃₂ ; 8,16-DiMeC ₃₂	478.55	3257	3263 [96]	378/379, 126/127, 308/309, 196/197; 378/379, 126/127, 280/281, 224/225; 378/379, 126/127, 252/253 (s)
B30	4-MeC ₃₂	464.53	3267	3265 [97]	420/421, 70/71
B31	6,16-DiMeC ₃₂	478.55	3285		406/407, 98/99, 252/253
B32	9-; 11-MeC ₃₃	478.55	3327	3335 [93]	364/365, 140/141; 336/337, 168/169
B33	13-; 15-; 17-MeC ₃₃	478.55	3330	3335 [93]	308/309, 196/197; 280/281, 224/225; 252/253 (s)
B34	9,23-DiMeC ₃₃	492.56	3359		378/379, 140/141, 350/351, 168/169

MM: molecular mass; KI: Kovats retention index; Ref.KI (Ref): reference KI if available with references in parenthesis; (s) indicates ions from symmetrical structures.



Figure 4. Internal standard-normalized peak areas of individual cuticular hydrocarbons (the alkanes in section B in Figure 2) in the two sexes of *B. tryoni*. Standardized peak areas were obtained by dividing the peak area of a compound by the peak area of the *n*-hexadecane internal standard. **B28** and **B29** co-eluted and hence the sum of their amounts are presented together. Error bars represent standard errors. The results of *t*-test comparisons between the sexes are shown (ns is not significant; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$).

3. Discussion

The present study finds that the *n*-hexane-extracted cuticular chemistry of *B. tryoni* includes a complex mixture of at least 66 compounds, including two spiroacetals, four aliphatic amides, 26 saturated/unsaturated C₁₂ to C₂₀ methyl, ethyl and propyl esters and 34 methyl branched saturated alkanes with a range of C₂₉ to C₃₃ carbon backbones. A previous study reported 14 cuticular compounds in *B. tryoni*, including five fatty acid esters, two siloxanes and seven methyl branched alkanes [98]. We did not detect the siloxanes or the methyl branched alkanes, which are all shorter than the alkanes found in the present study. We suspect that occurrence of the siloxanes in the previous work may reflect impurities, and incorrect assignments may have been given for the shorter methyl branched alkanes. The differences might have been also caused by technical differences; for example the previous work extracted cuticular compounds in methanol for 20 min [98], while the present study extracted in *n*-hexane for 3 min. The solvent choice of *n*-hexane was based on anticipated polarities of insect cuticular compounds that are generally less or non-polar [99]. The method used in the present study is more similar to methods widely used in studies of the cuticular hydrocarbons of other tephritids [53–59,100].

We found sexual dimorphism in several aspects of the cuticular chemistry of *B. tryoni*. In particular, our data suggest that spiroacetals and esters are specific to mature females. Sex-related differences in cuticular chemistry have also been reported in *B. dorsalis*, in which mature males have 7-monoenes that are absent from mature females and immatures of both sexes [100]. We also found quantitative differences between the sexes in their cuticular amides, which overall were more abundant in mature females than mature males, even though they are known to be particularly abundant in the rectal glands of mature males. Although the amides from male rectal glands have been suggested to function as male sex pheromones [65,66,68], the functions of these compounds on the cuticle, and in rectal glands of mature females, are not known. Our results indicate that more work is now needed on their functions in both sexes, both in rectal glands and on the cuticle.

Most of the non-alkane components which we find in cuticular extracts of *B. tryoni* have also been reported previously in rectal gland extracts of this species, with some of the same sex differences also evident [65,67–69]. In particular, the spiroacetals, amides and saturated/unsaturated esters we found in female cuticles had also been reported in female rectal glands. The presence of minute amounts of the amides and absence of the spiroacetals found on male cuticles also matches the earlier findings for male rectal glands, although the previous work also reported shorter chain esters (C₅ or less) in male rectal glands that were not found on cuticles. Notwithstanding the difference in the esters, the similarities between the two extract types otherwise suggest that at least some of the non-alkane components on the cuticle could originate in the rectal glands and be distributed over the body when grooming, as has been described in some social insects [101,102]. The differences in the esters and the quantitative differences in the amides noted above might in part reflect differences in volatility of the various compounds.

The C₁₂ to C₂₀ esters we found in cuticle of mature female *B. tryoni* have also been reported in the cuticle of females in five *B. dorsalis* complex species, but at different relative abundances between the species, suggesting that they may have a role in species recognition [58,59]. On the other hand, the methyl branched fatty acid esters found in the present study have not been previously reported in either the emission profile or rectal gland extracts of *B. tryoni* [67], or other fruit flies. Methyl branched fatty acid methyl, propyl or longer alkyl esters occur in other arthropods, such as spiders [103,104], but, to the best of our knowledge, specific roles have not yet been identified for them in any species. Branched fatty acid esters are biosynthetically feasible [105,106] and fatty acids are also the precursors of other cuticular compounds and hydrocarbons [2]. Hence, the branched esters found in this study may be intermediate products in the biosynthesis of other cuticular compounds.

While most of the C₂₉ to C₃₃ methyl branched alkanes detected in the present study have also been reported in other organisms (see references in Table 2), several of them have not previously been described in tephritids. Branched alkanes and/or unsaturated hydrocarbons have lower melting points than the corresponding normal alkanes, allowing the waxy layer of insect cuticles to be flexible over a

wide range of ambient temperatures [107,108]. Long chain hydrocarbons, and in particular branched and unsaturated alkanes, have also been linked to desiccation resistance in a variety of insects [109–111]. However, to what extent that applies in *B. tryoni* adults remains unclear. While qualitatively similar across the sexes and age groups studied, we found higher amounts of alkanes (as estimated from internal-standard-normalised peak areas) in mature than immature flies but, at least in domesticated flies from colonies maintained on artificial diets, desiccation resistance of *B. tryoni* decreases with age [73].

Several normal and methyl branched C₂₈ to C₄₀ alkanes have also been reported previously in two closely related taxa in the *Bactrocera dorsalis* complex, but as different isomers and in different amounts between the two [57]. Such variation in cuticular hydrocarbon chemistry has also been reported among species of another tephritid genus, *Anastrepha* [53,55,56,58,100], and even among populations within *Ceratitis rosa* [55]. As with the esters above, these taxon-specific cuticular hydrocarbon signatures may play a role in taxon recognition in nature, but they may also make excellent tools for taxonomists [53,55–58,100]. Their use in taxonomy is directly relevant to the *B. tryoni* complex which, despite its importance as a pest, is not well understood taxonomically. The *B. tryoni* complex includes three other taxa; *B. neohumeralis*, *B. aquilonis* and *B. melas* [60]. *Bactrocera tryoni* is clearly differentiated from *B. neohumeralis* in the timing of mating behavior [112], but the species status of *B. aquilonis* and *B. melas* is still debated [113] and the four taxa differ in their pest status and quarantine restrictions in various jurisdictions [113]. However, before cuticular chemistry can be used to resolve such taxonomic issues it will be important to investigate the extent to which it can be influenced by diet, maturity and the physical environment [114–116]. Given the population differences found in *Ceratitis rosa* [55], it will also be important to determine whether cuticular chemistry varies between geographic regions and during the course of domestication of species in the *B. tryoni* complex.

In summary, the present study provides the first detailed description of the cuticular chemistry of *B. tryoni*, and finds some clear qualitative and quantitative differences as flies mature and between the sexes. Our findings provide a foundation for studies addressing the roles of cuticular chemistry in functions such as desiccation resistance, protection from pathogens and injury, and chemical communication, as well as its potential application in resolving the taxonomy of the *B. tryoni* complex.

4. Materials and Methods

4.1. Chemicals

Ethyl dodecanoate (ethyl laurate), ethyl tridecanoate, propyl dodecanoate, methyl tetradecanoate, ethyl (*E*)-9-tetradecenoate (ethyl myristelaidate), ethyl (*Z*)-9-tetradecenoate (ethyl myristoleate), ethyl tetradecanoate (ethyl myristate), ethyl pentadecanoate, methyl (*Z*)-9-hexadecenoate, methyl hexadecanoate, ethyl (*E*)-9-hexadecenoate (ethyl palmitolelaidate) (*Z*)-9-hexadecenoate (ethyl palmitoleate), ethyl hexadecanoate (ethyl palmitate), ethyl (*Z,Z*)-octadeca-9,12-dienoate (ethyl linoleate), ethyl (*Z*)-9-octadecenoate (ethyl oleate), ethyl (*E*)-9-octadecenoate (ethyl elaidate), ethyl octadecanoate, hexadecane and straight-chain C₈–C₄₀ alkane standards were purchased from Sigma-Aldrich (St. Louis, MO, USA), Nu-Chek-Prep and INC (Minneapolis, MN, USA) or Alfa-Aesar. *N*-(3-methylbutyl)acetamide, *N*-(2-methylbutyl)propanamide, *N*-(3-methylbutyl)propanamide, *N*-(3-methylbutyl)isobutyramide were synthesized by reactions of an appropriate amine and an acid anhydride in water. Synthetic details are presented in Supplementary Materials. 2,8-Dimethyl-1,7-dioxaspiro[5.5]undecane was kindly provided by Ms. Sally Noushini.

4.2. Origin of Flies

Bactrocera tryoni larvae were collected from infested loquat fruits from a tree located in Marsfield NSW Australia (33.766080 S, 151.100722 E). A 500 mL plastic container with approximately 300 g of infested loquat fruits was placed on approximately a 1 cm deep layer of vermiculite in a 12.5 L plastic box for larvae to complete development and exit the fruit to pupate. Pupae were sieved and moved to

a fine mesh cage (47.5 × 47.5 × 47.5 cm) (Megaview Bugdorm 4S4545, Taiwan) where the flies emerged. Identity of *B. tryoni* was confirmed by examining emerged flies under a stereomicroscope using the key to tropical tephritid fruit flies [117] and Jane Royer kindly provided additional confirmation for the identity of *B. tryoni*. Adults were separated by sex within three days of emergence, when still sexually immature [118], and thereafter kept as single-sex cohorts of 80 flies in mesh cages (30.5 × 30.5 × 30.5 cm) (Megaview Bugdorm 4S4545, Taiwan). Adult flies were provided with food by coating a small area of the top of mesh cage with a paste prepared by mixing 15 g sugar, 5 g hydrolysate yeast (MP Biomedicals LLC) and 4 mL tap water. Water was provided by placing an inverted vial (6 cm height, 4 cm diameter) full of water on a sponge on the top of the mesh cage. All cages were maintained in a controlled environment room at 25 ± 0.5 °C, 65 ± 5% RH and photoperiod of 11.5:0.5:11.5:0.5 light: dusk: dark: dawn.

4.3. Extraction of Cuticular Compounds

Fifteen two-day old sexually immature flies of each sex and fifteen 20-day old sexually mature virgin flies of each sex were used for the extraction of cuticular compounds. The flies were killed by placing them in a 5 mL plastic vial on dry ice. Frozen flies were allowed to defrost at room temperature for three minutes immediately before the following extraction procedure. *n*-Hexane was chosen as the extraction solvent because many cuticular compounds found in other tephritids are less, or non-polar [53–59,100]. A single fly was immersed in 400 µL of *n*-hexane that contained 1.5 µg/mL *n*-hexadecane (Sigma-Aldrich, St. Louis, MO, USA) in a 1.5 mL clear glass vial. *n*-Hexadecane was used as an internal standard to normalize peak areas for comparisons between groups. The vial containing a fly was allowed to stand for three minutes at room temperature, then the fly was removed from the *n*-hexane extract. If *n*-hexane extracts contained aqueous droplets, the droplets were removed by adding sodium sulfate (Na₂SO₄) (Sigma-Aldrich, St. Louis, MO, USA) and gravity filtration. If samples contained solid organic matter, the solid was removed by gravity filtration. Gravity filtration was achieved by filtering the sample through a glass wool plug on the neck of a Pasteur pipette. *n*-Hexane extracts were concentrated to 120 µL under a gentle stream of nitrogen gas and transferred to a glass insert (150 µL) in a clear 1.5 mL GC vial. The samples were stored at 4 °C until analyzed.

4.4. Gas Chromatography Mass Spectrometry (GC-MS) Analysis

GC-MS analysis was carried out on a Shimadzu GCMS TQ8040 spectrometer equipped with a split/splitless injector and SH Rtx-5MS (30 m × 0.25 mm, 0.25 µm film) fused silica capillary column. The carrier gas was helium (99.999%) (BOC, North Ryde, NSW, Australia) at a flow rate of 1.0 mL/min. An aliquot of 1 µL sample was injected at splitless mode where the injector temperature was 270 °C. The temperature program was set initially at 50 °C (1 min), increased to 280 °C at a rate of 10 °C/min, then increased to 302 °C at a rate of 2 °C/min. The ion source and transfer line temperatures were 200 and 290 °C, respectively. The ionization method was electron impact at a voltage of 70 eV. The spectra were obtained over a mass range of *m/z* 47–650. The data were analyzed by Shimadzu GCMS Postrun program (Shimadzu, Kyoto, Japan) and compared with the mass fragmentation patterns in NIST libraries (NIST17-1, NIST17-2), and authentic standards. Retention indices (KI) were calculated with the Kovats retention index equation [119] and compared with KI values published in the literature. The structures of methyl-branched alkanes were assigned by using methods described in previous studies [120–122]. Briefly, the chain length and number of methyl groups of a methyl-branched alkane were established by examining an equivalent chain length and molar mass. The position of a methyl group was then assigned by examining fragment ions.

4.5. Comparison between Groups

The data were normalized by dividing the GC peak areas of individual components by the peak areas of the internal standard. The data were not normally distributed and hence were log-transformed for statistical analyses. However, the raw data were used to generate the graphs. The data for normal

and branched alkanes were analyzed by ANOVA to test for effects of sex and maturity on the amounts of compounds. The individual amides and alkanes were compared between sexes by *t*-test.

Supplementary Materials: The following are available online. Synthetic details of the amides and Figure S1. Mass spectra of 2-ethyl-8-methyl-1,7-dioxaspiro[5.5]undecane and methyl branched fatty acid esters.

Author Contributions: Conceptualization, V.M., P.W.T. and S.J.P.; methodology, V.M., P.W.T. and S.J.P.; formal analysis, G.P. and S.J.P.; investigation, C.C.-V., G.P., V.M. and S.J.P.; resources, P.W.T. and J.G.O.; writing—original draft preparation, S.J.P.; writing—review and editing, C.C.-V., G.P., J.G.O., P.W.T., V.M. and S.J.P.; supervision, J.G.O., P.W.T. and V.M.; project administration, P.W.T.; funding acquisition, P.W.T. All authors have read and agreed to the published version of the manuscript.

Funding: This research was conducted as part of the SITplus collaborative fruit fly program. Project *Raising Q-fly Sterile Insect Technique to World Standard* (HG14033) is funded by the Hort Frontiers Fruit Fly Fund, part of the Hort Frontiers strategic partnership initiative developed by Hort Innovation, with co-investment from Macquarie University and contributions from the Australian Government. This work received additional support from the Australian Research Council Industrial Transformation Training Centre (ITTC) for Fruit Fly Biosecurity Innovation (Project IC50100026), funded by the Australian Government, including a fellowship to SJP.

Acknowledgments: We thank Sally Noushini for providing 2,8-dimethyl-1,7-dioxaspiro[5,5]undecane, and Jane Royer for expert identification of *B. tryoni*.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Gibbs, A.; Crockett, E.L. The Biology of Lipids: Integrative and Comparative Perspectives. *Am. Zool.* **1998**, *38*, 265–267. [[CrossRef](#)]
2. Howard, R.W.; Blomquist, G.J. Chemical Ecology and Biochemistry of Insect Hydrocarbons. *Annu. Rev. Entomol.* **1982**, *27*, 149–172. [[CrossRef](#)]
3. Lockey, K.H. Lipids of the insect cuticle: Origin, composition and function. *Comp. Biochem. Physiol. Part B Comp. Biochem.* **1988**, *89*, 595–645. [[CrossRef](#)]
4. Blomquist, G.J.; Jackson, L.L. Chemistry and biochemistry of insect waxes. *Prog. Lipid Res.* **1979**, *17*, 319–345. [[CrossRef](#)]
5. Blomquist, G.J.; Nelson, D.R.; De Renobales, M. Chemistry, biochemistry, and physiology of insect cuticular lipids. *Arch. Insect Biochem. Physiol.* **1987**, *6*, 227–265. [[CrossRef](#)]
6. Ferveur, J.-F. Cuticular Hydrocarbons: Their Evolution and Roles in Drosophila Pheromonal Communication. *Behav. Genet.* **2005**, *35*, 279–295. [[CrossRef](#)]
7. Gołębiowski, M.; Boguś, M.I.; Paszkiewicz, M.; Stepnowski, P. Cuticular lipids of insects as potential biofungicides: Methods of lipid composition analysis. *Anal. Bioanal. Chem.* **2010**, *399*, 3177–3191. [[CrossRef](#)]
8. Gołębiowski, M.; Urbanek, A.; Pietrzak, A.; Naczka, A.M.; Bojke, A.; Tkaczuk, C.; Stepnowski, P. Effects of the entomopathogenic fungus *Metarhizium flavoviride* on the fat body lipid composition of *Zophobas morio* larvae (Coleoptera: Tenebrionidae). *Naturwissenschaften* **2020**, *107*, 1–11. [[CrossRef](#)]
9. Gołębiowski, M.; Cerkowniak, M.; Urbanek, A.; Dawgul, M.; Kamysz, W.; Boguś, M.I.; Stepnowski, P. Identification and antifungal activity of novel organic compounds found in cuticular and internal lipids of medically important flies. *Microbiol. Res.* **2015**, *170*, 213–222. [[CrossRef](#)]
10. Boguś, M.I.; Czygier, M.; Gołębiowski, M.; Kędra, E.; Kucińska, J.; Mazgajska, J.; Samborski, J.; Wieloch, W.; Włóka, E. Effects of insect cuticular fatty acids on in vitro growth and pathogenicity of the entomopathogenic fungus *Conidiobolus coronatus*. *Exp. Parasitol.* **2010**, *125*, 400–408. [[CrossRef](#)]
11. Urbanek, A.; Szadziewski, R.; Stepnowski, P.; Boros-Majewska, J.; Gabriel, I.; Dawgul, M.; Kamysz, W.; Sosnowska, D.; Gołębiowski, M. Composition and antimicrobial activity of fatty acids detected in the hygroscopic secretion collected from the secretory setae of larvae of the biting midge *Forcipomyia nigra* (Diptera: Ceratopogonidae). *J. Insect Physiol.* **2012**, *58*, 1265–1276. [[CrossRef](#)] [[PubMed](#)]
12. Gołębiowski, M.; Cerkowniak, M.; Dawgul, M.; Kamysz, W.; Boguś, M.I.; Stepnowski, P. The antifungal activity of the cuticular and internal fatty acid methyl esters and alcohols in *Calliphora vomitoria*. *Parasitology* **2013**, *140*, 972–985. [[CrossRef](#)]

13. Howard, R.W.; Blomquist, G.J. Ecological, behavioral, and biochemical aspects of insect hydrocarbons. *Annu. Rev. Entomol.* **2005**, *50*, 371–393. [[CrossRef](#)] [[PubMed](#)]
14. Carlson, D.A.; Nelson, D.R.; Langley, P.A.; Coates, T.W.; Davis, T.L.; Der Linden, M.E.L.-V. Contact sex pheromone in the tsetse fly *Glossina pallidipes* (Austen) Identification and Synthesis. *J. Chem. Ecol.* **1984**, *10*, 429–450. [[CrossRef](#)]
15. Ginzel, M.D.; Blomquist, G.J.; Millar, J.G.; Hanks, L.M. Role of contact pheromones in mate recognition in *Xylotrechus colonus*. *J. Chem. Ecol.* **2003**, *29*, 533–545. [[CrossRef](#)]
16. Wilson, E.O. *Chemical Communication within Animal Species*; Academic Press, Inc.: New York, NY, USA, 1970; pp. 133–155.
17. Xue, H.-J.; Segraves, K.A.; Wei, J.; Zhang, B.; Nie, R.-E.; Li, W.-Z.; Yang, X.-K. Chemically mediated sexual signals restrict hybrid speciation in a flea beetle. *Behav. Ecol.* **2018**, *29*, 1462–1471. [[CrossRef](#)]
18. Dettner, K.; Liepert, C. Chemical Mimicry and Camouflage. *Annu. Rev. Entomol.* **1994**, *39*, 129–154. [[CrossRef](#)]
19. Heifetz, Y.; Voet, H.; Applebaum, S.W. Factors affecting behavioral phase transition in the desert locust, *Schistocerca gregaria* (Forskål) (Orthoptera: Acrididae). *J. Chem. Ecol.* **1996**, *22*, 1717–1734. [[CrossRef](#)]
20. Carlson, D.A.; Mayer, M.S.; Silhacek, D.L.; James, J.D.; Beroza, M.; Bierl, B.A. Sex Attractant Pheromone of the House Fly: Isolation, Identification and Synthesis. *Science* **1971**, *174*, 76–78. [[CrossRef](#)]
21. Jennings, J.H.; Etges, W.J.; Schmitt, T.; Hoikkala, A. Cuticular hydrocarbons of *Drosophila montana*: Geographic variation, sexual dimorphism and potential roles as pheromones. *J. Insect Physiol.* **2014**, *61*, 16–24. [[CrossRef](#)]
22. Roelofs, W.L.; Liu, W.; Hao, G.; Jiao, H.; Rooney, A.P.; Linn, J.C.E. Evolution of moth sex pheromones via ancestral genes. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 13621–13626. [[CrossRef](#)] [[PubMed](#)]
23. Mant, J.; Brändli, C.; Vereecken, N.J.; Schulz, C.M.; Francke, W.; Schiestl, F.P. Cuticular Hydrocarbons as Sex Pheromone of the Bee *Colletes cunicularius* and the Key to its Mimicry by the Sexually Deceptive Orchid, *Ophrys exaltata*. *J. Chem. Ecol.* **2005**, *31*, 1765–1787. [[CrossRef](#)] [[PubMed](#)]
24. Nojima, S.; Shimomura, K.; Honda, H.; Yamamoto, I.; Ohsawa, K. Contact Sex Pheromone Components of the Cowpea Weevil, *Callosobruchus maculatus*. *J. Chem. Ecol.* **2007**, *33*, 923–933. [[CrossRef](#)]
25. Keeseey, I.W.; Koerte, S.; Retzke, T.; Haverkamp, A.; Hansson, B.S.; Knaden, M. Adult Frass Provides a Pheromone Signature for *Drosophila* Feeding and Aggregation. *J. Chem. Ecol.* **2016**, *42*, 739–747. [[CrossRef](#)]
26. Mitaka, Y.; Matsuyama, S.; Mizumoto, N.; Matsuura, K.; Akino, T. Chemical identification of an aggregation pheromone in the termite *Reticulitermes speratus*. *Sci. Rep.* **2020**, *10*, 7424. [[CrossRef](#)]
27. Hamilton, J.A.; Wada-Katsumata, A.; Schal, C. Role of Cuticular Hydrocarbons in German Cockroach (Blattodea: Ectobiidae) Aggregation Behavior. *Environ. Entomol.* **2019**, *48*, 546–553. [[CrossRef](#)] [[PubMed](#)]
28. Cooper, W.E.; Andersson, M. Sexual Selection. *Copeia* **1995**, 756. [[CrossRef](#)]
29. Blomquist, G.J.; Bagnères, A.-G. Introduction: History and overview of insect hydrocarbons. In *Insect Hydrocarbons: Biology, Biochemistry, and Chemical Ecology*; Bagnères, A.-G., Blomquist, G.J., Eds.; Cambridge University Press: Cambridge, UK, 2010; pp. 3–18.
30. Luo, Y.; Zhang, Y.; Farine, J.; Ferveur, J.; Ramírez, S.; Kopp, A. Evolution of sexually dimorphic pheromone profiles coincides with increased number of male-specific chemosensory organs in *Drosophila prolongata*. *Ecol. Evol.* **2019**, *9*, 13608–13618. [[CrossRef](#)]
31. Ngumbi, E.N.; Hanks, L.M.; Suarez, A.V.; Millar, J.G.; Berenbaum, M.R. Author Correction: Factors Associated with Variation in Cuticular Hydrocarbon Profiles in the Navel Orangeworm, *Amyelois transitella* (Lepidoptera: Pyralidae). *J. Chem. Ecol.* **2020**, *46*, 232. [[CrossRef](#)]
32. Farine, J.-P.; Ferveur, J.-F.; Everaerts, C. Volatile *Drosophila* Cuticular Pheromones Are Affected by Social but Not Sexual Experience. *PLoS ONE* **2012**, *7*, e40396. [[CrossRef](#)]
33. Dias, N.P.; Zotti, M.J.; Montoya, P.; Carvalho, I.R.; Nava, D.E. Fruit fly management research: A systematic review of monitoring and control tactics in the world. *Crop. Prot.* **2018**, *112*, 187–200. [[CrossRef](#)]
34. Robledo, N.; Arzuffi, R.; Robledo-Quintos, N. Influence of Host Fruit and Conspecifics on the Release of the Sex Pheromone By *Toxotrypana curvicauda* Males (Diptera: Tephritidae). *Environ. Entomol.* **2012**, *41*, 387–391. [[CrossRef](#)] [[PubMed](#)]
35. Tan, H.K.; Nishida, R. Incorporation of Raspberry Ketone in the Rectal Glands of Males of the Queensland Fruit Fly, *Bactrocera tryoni* FROGGATT (Diptera: Tephritidae). *Appl. Entomol. Zool.* **1995**, *30*, 494–497. [[CrossRef](#)]

36. Jang, E.B.; Carvalho, L.A.F.N.; Chen, C.-C.; Siderhurst, M.S. Cucumber Lure Trapping of *Zeugodacus cucurbitae* (Diptera:Tephritidae) in Hawaii and Taiwan: Longevity and Nontargets Captures. *J. Econ. Entomol.* **2016**, *110*, 201–207. [[CrossRef](#)]
37. Royer, J.E.; De Faveri, S.G.; Lowe, G.E.; Wright, C.L. Cucumber volatile blend, a promising female-biased lure for *Bactrocera cucumis* (French 1907) (Diptera: Tephritidae: Dacinae), a pest fruit fly that does not respond to male attractants. *Austral Entomol.* **2014**, *53*, 347–352. [[CrossRef](#)]
38. Siderhurst, M.S.; Jang, E.B. Female-Biased Attraction of Oriental Fruit Fly, *Bactrocera dorsalis* (Hendel), to a Blend of Host Fruit Volatiles From *Terminalia catappa* L. *J. Chem. Ecol.* **2006**, *32*, 2513–2524. [[CrossRef](#)]
39. Siderhurst, M.S.; Jang, E.B. Attraction of female Oriental fruit fly, *Bactrocera dorsalis*, to *Terminalia catappa* fruit in wind tunnel and olfactometer tests. *Formos. Entomol.* **2006**, *26*, 45–55.
40. Siderhurst, M.S.; Jang, E.B. Cucumber Volatile Blend Attractive to Female Melon Fly, *Bactrocera cucurbitae* (Coquillett). *J. Chem. Ecol.* **2010**, *36*, 699–708. [[CrossRef](#)]
41. Tan, K.H. Fruit fly pests as pollinators of wild orchids. In Proceedings of the 7th International Symposium on Fruit Flies of Economic Importance, Salvador, Brazil, 10–15 September 2006; pp. 195–206.
42. Tan, K.H.; Nishida, R. Zingerone in the floral synomone of *Bulbophyllum baileyi* (Orchidaceae) attracts *Bactrocera* fruit flies during pollination. *Biochem. Syst. Ecol.* **2007**, *35*, 334–341. [[CrossRef](#)]
43. Tan, K.H.; Tan, L.T.; Nishida, R. Floral Phenylpropanoid Cocktail and Architecture of *Bulbophyllum vinaceum* Orchid in Attracting Fruit Flies for Pollination. *J. Chem. Ecol.* **2006**, *32*, 2429–2441. [[CrossRef](#)]
44. Tan, K.-H.; Nishida, R. Synomone or kairomone?—*Bulbophyllum apertum* flower releases raspberry ketone to attract *Bactrocera* fruit flies. *J. Chem. Ecol.* **2005**, *31*, 497–507.
45. Tan, K.-H.; Nishida, R.; Toong, Y.-C. Floral synomone of a wild orchid, *Bulbophyllum cheiri*, lures *Bactrocera* fruit flies for pollination. *J. Chem. Ecol.* **2002**, *28*, 1161–1172. [[CrossRef](#)] [[PubMed](#)]
46. Jang, E.B.; Carvalho, L.A.; Stark, J.D. Attraction of Female Oriental Fruit Fly, *Bactrocera dorsalis*, to Volatile Semiochemicals from Leaves and Extracts of a Nonhost Plant, Panax (*Polyscias guilfoylei*) in Laboratory and Olfactometer Assays. *J. Chem. Ecol.* **1997**, *23*, 1389–1401. [[CrossRef](#)]
47. Siderhurst, M.S.; Park, S.J.; Buller, C.N.; Jamie, I.M.; Manoukis, N.C.; Jang, E.B.; Taylor, P.W. Raspberry Ketone Trifluoroacetate, a New Attractant for the Queensland Fruit Fly, *Bactrocera tryoni* (Froggatt). *J. Chem. Ecol.* **2016**, *42*, 156–162. [[CrossRef](#)]
48. Kather, R.; Martin, S.J. Cuticular hydrocarbon profiles as a taxonomic tool: Advantages, limitations and technical aspects. *Physiol. Entomol.* **2012**, *37*, 25–32. [[CrossRef](#)]
49. Vanickova, L.; Hernández-Ortiz, V.; Bravo, I.S.J.; Dias, V.S.; Roriz, A.K.P.; Laumann, R.A.; Mendonça, A.D.L.; Paranhos, B.A.J.; Nascimento, R.R.D. Current knowledge of the species complex *Anastrepha fraterculus* (Diptera, Tephritidae) in Brazil. *ZooKeys* **2015**, *540*, 211–237. [[CrossRef](#)]
50. Carlson, D.A.; Yocom, S.R. Cuticular hydrocarbons from six species of tephritid fruit flies. *Arch. Insect Biochem. Physiol.* **1986**, *3*, 397–412. [[CrossRef](#)]
51. Sutton, B.D.; Carlson, D.A. Interspecific variation in tephritid fruit fly larvae surface hydrocarbons. *Arch. Insect Biochem. Physiol.* **1993**, *23*, 53–65. [[CrossRef](#)]
52. Sutton, B.D.; Steck, G.J. Discrimination of Caribbean and Mediterranean Fruit Fly Larvae (Diptera: Tephritidae) by Cuticular Hydrocarbon Analysis. *Fla. Entomol.* **1994**, *77*, 231. [[CrossRef](#)]
53. Vaníčková, L.; Virgilio, M.; Tomčala, A.; Břízová, R.; Ekesi, S.; Hoskovec, M.; Kalinová, B.; Do Nascimento, R.R.; De Meyer, M. Resolution of three cryptic agricultural pests (*Ceratitis fasciventris*, *C. anonae*, *C. rosa*, Diptera: Tephritidae) using cuticular hydrocarbon profiling. *Bull. Entomol. Res.* **2014**, *104*, 631–638.
54. Vanickova, L.; Břízová, R.; Mendonça, A.L.; Pompeiano, A.; Nascimento, R.R.D. Intraspecific variation of cuticular hydrocarbon profiles in the *Anastrepha fraterculus* (Diptera:Tephritidae) species complex. *J. Appl. Entomol.* **2015**, *139*, 679–689. [[CrossRef](#)]
55. Vanickova, L.; Břízová, R.; Pompeiano, A.; Ekesi, S.; De Meyer, M. Cuticular hydrocarbons corroborate the distinction between lowland and highland Natal fruit fly (Tephritidae, *Ceratitis rosa*) populations. *ZooKeys* **2015**, *540*, 507–524. [[CrossRef](#)] [[PubMed](#)]
56. Vanickova, L.; Břízová, R.; Pompeiano, A.; Ferreira, L.L.; De Aquino, N.C.; Tavares, R.D.F.; Rodriguez, L.D.; Mendonça, A.D.L.; Canal, N.A.; Nascimento, R.R.D. Characterisation of the chemical profiles of Brazilian and Andean morphotypes belonging to the *Anastrepha fraterculus* complex (Diptera, Tephritidae). *ZooKeys* **2015**, *540*, 193–209. [[CrossRef](#)] [[PubMed](#)]

57. Goh, S.; Ooi, K.; Chuah, C.; Yong, H.; Khoo, S.; Ong, S. Cuticular hydrocarbons from two species of Malaysian *Bactrocera* fruit flies. *Biochem. Syst. Ecol.* **1993**, *21*, 215–226. [[CrossRef](#)]
58. Vanickova, L.; Nagy, R.; Pompeiano, A.; Kalinová, B. Epicuticular chemistry reinforces the new taxonomic classification of the *Bactrocera dorsalis* species complex (Diptera: Tephritidae, Dacinae). *PLoS ONE* **2017**, *12*, e0184102. [[CrossRef](#)]
59. Shen, J.; Hu, L.; Zhou, X.; Dai, J.; Chen, B.; Li, S. Allyl-2,6-dimethoxyphenol, a female-biased compound, is robustly attractive to conspecific males of *Bactrocera dorsalis* at close range. *Entomol. Exp. Appl.* **2019**, *167*, 811–819. [[CrossRef](#)]
60. Clarke, A.R.; Powell, K.S.; Weldon, C.W.; Taylor, P.W. The ecology of *Bactrocera tryoni* (Diptera: Tephritidae): What do we know to assist pest management? *Ann. Appl. Biol.* **2011**, *158*, 26–54. [[CrossRef](#)]
61. Gilchrist, A.S.; Dominiak, B.; Gillespie, P.S.; Sved, J.A. Variation in population structure across the ecological range of the Queensland fruit fly. *Bactrocera tryoni*. *Aust. J. Zool.* **2006**, *54*, 87. [[CrossRef](#)]
62. Sutherst, R.W.; Collyer, B.S.; Yonow, T. The vulnerability of Australian horticulture to the Queensland fruit fly, *Bactrocera* (*Dacus*) *tryoni*, under climate change. *Aust. J. Agric. Res.* **2000**, *51*, 467. [[CrossRef](#)]
63. Dominiak, B.; Ekman, J.H.; Broughton, S. Mass trapping and other management options for Mediterranean fruit fly and Queensland fruit fly in Australia. *Gen. App. Entomol.* **2016**, *44*, 1–8.
64. Florec, V.; Sadler, R.J.; White, B.; Dominiak, B.C.; White, B. Choosing the battles: The economics of area wide pest management for Queensland fruit fly. *Food Policy* **2013**, *38*, 203–213. [[CrossRef](#)]
65. Bellas, T.E.; Fletcher, B.S. Identification of the major components in the secretion from the rectal pheromone glands of the Queensland fruit flies *Dacus tryoni* and *Dacus neohumeralis* (Diptera: Tephritidae). *J. Chem. Ecol.* **1979**, *5*, 795–803. [[CrossRef](#)]
66. Fletcher, B.S. Storage and Release of a Sex Pheromone by the Queensland Fruit Fly, *Dacus tryoni* (Diptera: Trypetidae). *Nature* **1968**, *219*, 631–632. [[CrossRef](#)] [[PubMed](#)]
67. El-Sayed, A.M.; Venkatesham, U.; Unelius, C.R.; Sporle, A.; Pérez, J.; Taylor, P.W.; Suckling, D.M. Chemical Composition of the Rectal Gland and Volatiles Released by Female Queensland Fruit Fly, *Bactrocera tryoni* (Diptera: Tephritidae). *Environ. Entomol.* **2019**, *48*, 807–814. [[CrossRef](#)]
68. Pérez, J.; Park, S.J.; Taylor, P.W. Domestication modifies the volatile emissions produced by male Queensland fruit flies during sexual advertisement. *Sci. Rep.* **2018**, *8*, 16503. [[CrossRef](#)] [[PubMed](#)]
69. Noushini, S.; Park, S.J.; Jamie, I.M.; Jamie, J.F.; Taylor, P.W. Sampling technique biases in the analysis of fruit fly pheromones: A case study of Queensland fruit fly. *Sci. Rep.* **2020**, in press.
70. Booth, Y.K.; Schwartz, B.D.; Fletcher, M.; Lambert, L.K.; Kitching, W.; De Voss, J.J. A diverse suite of spiroacetals, including a novel branched representative, is released by female *Bactrocera tryoni* (Queensland fruit fly). *Chem. Commun.* **2006**, *42*, 3975–3977. [[CrossRef](#)]
71. Lin, Y.; Jin, T.; Zeng, L.; Lu, Y. Cuticular penetration of β -cypermethrin in insecticide-susceptible and resistant strains of *Bactrocera dorsalis*. *Pestic. Biochem. Physiol.* **2012**, *103*, 189–193. [[CrossRef](#)]
72. Sultana, S.; Baumgartner, J.B.; Dominiak, B.C.; Royer, J.E.; Beaumont, L.J. Potential impacts of climate change on habitat suitability for the Queensland fruit fly. *Sci. Rep.* **2017**, *7*, 13025. [[CrossRef](#)]
73. Weldon, C.W.; Taylor, P.W. Desiccation resistance of adult Queensland fruit flies *Bactrocera tryoni* decreases with age. *Physiol. Entomol.* **2010**, *35*, 385–390. [[CrossRef](#)]
74. Weldon, C.W.; Yap, S.; Taylor, P. Desiccation resistance of wild and mass-reared *Bactrocera tryoni* (Diptera: Tephritidae). *Bull. Entomol. Res.* **2013**, *103*, 690–699. [[CrossRef](#)] [[PubMed](#)]
75. Dominiak, B.C.; Westcott, A.E.; Barchia, I.M. Release of sterile Queensland fruit fly, *Bactrocera* > *tryoni* (Froggatt) (Diptera: Tephritidae), at Sydney, Australia. *Aust. J. Exp. Agric.* **2003**, *43*, 519–528. [[CrossRef](#)]
76. Meats, A.; Duthie, R.; Clift, A.D.; Dominiak, B.C. Trials on variants of the Sterile Insect Technique (SIT) for suppression of populations of the Queensland fruit fly in small towns neighbouring a quarantine (exclusion) zone. *Aust. J. Exp. Agric.* **2003**, *43*, 389–395. [[CrossRef](#)]
77. Reynolds, O.L.; Smallridge, C.J.; Cockington, V.G.; Penrose, L.D. The effect of release method and trial site on recapture rates of adult sterile Queensland fruit fly, *Bactrocera tryoni* (Froggatt) (Diptera: Tephritidae). *Aust. J. Entomol.* **2011**, *51*, 116–126. [[CrossRef](#)]
78. Noushini, S.; Pérez, J.; Park, S.J.; Holgate, D.; Alvarez, V.M.; Jamie, I.M.; Jamie, J.F.; Taylor, P.W. Attraction and Electrophysiological Response to Identified Rectal Gland Volatiles in *Bactrocera frauenfeldi* (Schiner). *Molecules* **2020**, *25*, 1275. [[CrossRef](#)]

79. El-Sayed, A.M.; Heppelthwaite, V.J.; Manning, L.-A.; Gibb, A.R.; Suckling, D.M.; El-Sayed, A.M. Volatile Constituents of Fermented Sugar Baits and Their Attraction to Lepidopteran Species. *J. Agric. Food Chem.* **2005**, *53*, 953–958. [[CrossRef](#)]
80. Pino, J.A.; Mesa, J.; Muñoz, Y.; Martí, M.P.; Marbot, R. Volatile Components from Mango (*Mangifera indica* L.) Cultivars. *J. Agric. Food Chem.* **2005**, *53*, 2213–2223. [[CrossRef](#)]
81. Peppard, T.L. Volatile flavor constituents of *Monstera deliciosa*. *J. Agric. Food Chem.* **1992**, *40*, 257–262. [[CrossRef](#)]
82. Silva, D.B.; Pott, A.; Oliveira, D.C.R. Analyses of the Headspace Volatile Constituents of Aerial Parts (leaves and stems), Flowers and Fruits of *Bidens gardneri* Bak. and *Bidens sulphurea* (Cav.) Sch.Bip. Using Solid-Phase Microextraction. *J. Essent. Oil Res.* **2010**, *22*, 560–563. [[CrossRef](#)]
83. Isidorov, V.; Krajewska, U.; Dubis, E.N.; Jdanova, M. Partition coefficients of alkyl aromatic hydrocarbons and esters in a hexane-acetonitrile system. *J. Chromatogr. A* **2001**, *923*, 127–136. [[CrossRef](#)]
84. Robinson, A.L.; Adams, D.O.; Boss, P.K.; Heymann, H.; Solomon, P.S.; Trengove, R. Influence of Geographic Origin on the Sensory Characteristics and Wine Composition of *Vitis vinifera* cv. Cabernet Sauvignon Wines from Australia. *Am. J. Enol. Vitic.* **2012**, *63*, 467–476. [[CrossRef](#)]
85. Palmeira, S.F.; Moura, F.D.S.; Alves, V.D.L.; De Oliveira, F.M.; Bento, E.S.; Conserva, L.M.; Andrade, E.D.A. Neutral components from hexane extracts of *Croton sellowii*. *Flavour Fragr. J.* **2004**, *19*, 69–71. [[CrossRef](#)]
86. Pino, J.A.; Almora, K.; Marbot, R. Volatile components of papaya (*Carica papaya* L., Maradol variety) fruit. *Flavour Fragr. J.* **2003**, *18*, 492–496. [[CrossRef](#)]
87. Palmeira, S.F.; Conserva, L.M.; Andrade, E.H.D.A.; Guilhon, G.M.S.P. Analysis by GC-MS of the hexane extract of the aerial parts of *Aristolochia acutifolia* Duchtr. *Flavour Fragr. J.* **2001**, *16*, 85–88. [[CrossRef](#)]
88. Demyttenaere, J.C.; Martínez, J.I.S.; Verhé, R.; Sandra, P.; De Kimpe, N.; Sanchezmartinez, J. Analysis of volatiles of malt whisky by solid-phase microextraction and stir bar sorptive extraction. *J. Chromatogr. A* **2003**, *985*, 221–232. [[CrossRef](#)]
89. Andriamaharavo, N.R. Retention Data: NIST Mass Spectrometry Data Center. 2014. Available online: <https://webbook.nist.gov/cgi/cbook.cgi?Source=2014AND%2319410M&Units=SI&Mask=2000> (accessed on 10 September 2020).
90. Tuberoso, C.; Barra, A.; Angioni, A.; Sarritzu, E.; Pirisi, F.M. Chemical Composition of Volatiles in Sardinian Myrtle (*Myrtus communis* L.) Alcoholic Extracts and Essential Oils. *J. Agric. Food Chem.* **2006**, *54*, 1420–1426. [[CrossRef](#)]
91. Kenig, F.; Damsté, J.S.S.; Dalen, A.K.-V.; Rijpstra, W.C.; Huc, A.Y.; De Leeuw, J.W. Occurrence and origin of mono-, di-, and trimethylalkanes in modern and Holocene cyanobacterial mats from Abu Dhabi, United Arab Emirates. *Geochim. Cosmochim. Acta* **1995**, *59*, 2999–3015. [[CrossRef](#)]
92. Nelson, D.R.; Charlet, L.D. Cuticular hydrocarbons of the sunflower beetle, *Zygogramma exclamationis*. *Comp. Biochem. Physiol. Part B Biochem. Mol. Boil.* **2003**, *135*, 273–284. [[CrossRef](#)]
93. Carlson, D.; Geden, C.; Bernier, U. Identification of Pupal Exuviae of *Nasonia vitripennis* and *Muscidifurax raptorellus* Parasitoids Using Cuticular Hydrocarbons. *Boil. Control.* **1999**, *15*, 97–106. [[CrossRef](#)]
94. Lockey, K.H. Cuticular hydrocarbons of adult *Cylindrinotus laevioctostriatus* (Goeze) and *Phylan gibbus* (Fabricius) (Coleoptera: Tenebrionidae). *Insect Biochem.* **1981**, *11*, 549–561. [[CrossRef](#)]
95. Katritzky, A.R.; Chen, K.; Maran, U.; Carlson, D.A. QSPR correlation and predictions of GC retention indexes for methyl-branched hydrocarbons produced by insects. *Anal. Chem.* **2000**, *72*, 101–109. [[CrossRef](#)] [[PubMed](#)]
96. Lockey, K.H. The adult cuticular hydrocarbons of *Tenebrio molitor* L. and *Tenebrio obscurus* F. (Coleoptera: Tenebrionidae). *Insect Biochem.* **1978**, *8*, 237–250. [[CrossRef](#)]
97. Bernier, U.R.; Carlson, D.A.; Geden, C.J. Gas chromatography/mass spectrometry analysis of the cuticular hydrocarbons from parasitic wasps of the genus *Muscidifurax*. *J. Am. Soc. Mass Spectrom.* **1998**, *9*, 320–332. [[CrossRef](#)]
98. Ekanayake, D. [*Leave for Adam-Compressed File*] *The Mating System and Courtship Behaviour of the Queensland Fruit Fly, Bactrocera tryoni* (Froggatt) (Diptera: Tephritidae); Queensland University of Technology: Brisbane, Australia, 2017; p. 271.
99. Blomquist, G.J.; Tittiger, C.; Jurenka, R. Cuticular Hydrocarbons and Pheromones of Arthropods. In *Hydrocarbons, Oils and Lipids: Diversity, Origin, Chemistry and Fate*. Available online: https://doi.org/10.1007/978-3-319-54529-5_11-1 (accessed on 10 September 2020).

100. Vanickova, L.; Svatoš, A.; Kroiss, J.; Kaltenpoth, M.; Nascimento, R.R.D.; Hoskovec, M.; Břízová, R.; Kalinová, B. Cuticular Hydrocarbons of the South American Fruit Fly *Anastrepha fraterculus*: Variability with Sex and Age. *J. Chem. Ecol.* **2012**, *38*, 1133–1142. [[CrossRef](#)]
101. Lucas, C.; Pho, D.; Fresneau, D.; Jallon, J. Hydrocarbon circulation and colonial signature in *Pachycondyla villosa*. *J. Insect Physiol.* **2004**, *50*, 595–607. [[CrossRef](#)]
102. Soroker, V.; Vienne, C.; Hefetz, A. Hydrocarbon dynamics within and between nestmates in *Cataglyphis niger* (Hymenoptera: Formicidae). *J. Chem. Ecol.* **1995**, *21*, 365–378. [[CrossRef](#)]
103. Bagnères, A.-G.; Tralalon, M.; Blomquist, G.J.; Schulz, S. Waxes of the social spider *Anelosimus eximius* (Araneae, Theridiidae): Abundance of novel n-propyl esters of long-chain methyl-branched fatty acids. *Arch. Insect Biochem. Physiol.* **1997**, *36*, 295–314. [[CrossRef](#)]
104. Chinta, S.P.; Goller, S.; Uhl, G.; Schulz, S. Identification and Synthesis of Branched Wax-type Esters, Novel Surface Lipids from the Spider *Argyrodes elevates* (Araneae: Theridiidae). *Chem. Biodivers.* **2016**, *13*, 1202–1220. [[CrossRef](#)]
105. Gu, P.; Welch, W.H.; Blomquist, G.J. Methyl-branched fatty acid biosynthesis in the German cockroach, *Blattella germanica*: Kinetic studies comparing a microsomal and soluble fatty acid synthetase. *Insect Biochem. Mol. Boil.* **1993**, *23*, 263–271. [[CrossRef](#)]
106. Wang, H.-L.; Brattström, O.; Brakefield, P.M.; Francke, W.; Löfstedt, C. Identification and Biosynthesis of Novel Male Specific Esters in the Wings of the Tropical Butterfly, *Bicyclus martius sanaos*. *J. Chem. Ecol.* **2014**, *40*, 549–559. [[CrossRef](#)]
107. Morgan, E. *Biosynthesis in Insects*; Royal Society of Chemistry (RSC): Cambridge, UK, 2006.
108. Gibbs, A.; Pomonis, J. Physical properties of insect cuticular hydrocarbons: The effects of chain length, methyl-branching and unsaturation. *Comp. Biochem. Physiol. Part B Biochem. Mol. Boil.* **1995**, *112*, 243–249. [[CrossRef](#)]
109. Foley, B.R.; Telonis-Scott, M. Quantitative genetic analysis suggests causal association between cuticular hydrocarbon composition and desiccation survival in *Drosophila melanogaster*. *Heredity* **2010**, *106*, 68–77. [[CrossRef](#)] [[PubMed](#)]
110. Gefen, E.; Talal, S.; Brendzel, O.; Dror, A.; Fishman, A. Variation in quantity and composition of cuticular hydrocarbons in the scorpion *Buthus occitanus* (Buthidae) in response to acute exposure to desiccation stress. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* **2015**, *182*, 58–63. [[CrossRef](#)] [[PubMed](#)]
111. Ferveur, J.-F.; Cortot, J.; Rihani, K.; Cobb, M.; Everaerts, C. Desiccation resistance: Effect of cuticular hydrocarbons and water content in *Drosophila melanogaster* adults. *PeerJ* **2018**, *6*, 4318. [[CrossRef](#)] [[PubMed](#)]
112. Pike, N.; Wang, W.Y.; Meats, A. The likely fate of hybrids of *Bactrocera tryoni* and *Bactrocera neohumeralis*. *Heredity* **2003**, *90*, 365–370. [[CrossRef](#)] [[PubMed](#)]
113. Popa-Báez, Á.-D.; Catullo, R.; Lee, S.F.; Yeap, H.L.; Mourant, R.G.; Frommer, M.; Sved, J.A.; Cameron, E.C.; Edwards, O.R.; Taylor, P.W.; et al. Genome-wide patterns of differentiation over space and time in the Queensland fruit fly. *Sci. Rep.* **2020**, *10*, 1–13. [[CrossRef](#)]
114. Fedina, T.Y.; Kuo, T.-H.; Dreisewerd, K.; Dierick, H.A.; Yew, J.Y.; Pletcher, S.D. Dietary Effects on Cuticular Hydrocarbons and Sexual Attractiveness in *Drosophila*. *PLoS ONE* **2012**, *7*, e49799. [[CrossRef](#)]
115. Otte, T.; Hilker, M.; Geiselhardt, S. The effect of dietary fatty acids on the cuticular hydrocarbon phenotype of an herbivorous insect and consequences for mate recognition. *J. Chem. Ecol.* **2015**, *41*, 32–43. [[CrossRef](#)]
116. Otte, T.; Hilker, M.; Geiselhardt, S. Phenotypic Plasticity of Cuticular Hydrocarbon Profiles in Insects. *J. Chem. Ecol.* **2018**, *44*, 235–247. [[CrossRef](#)]
117. Drew, R.A.I. The tropical fruit flies (Diptera: Tephritidae: Dacinae) of the Australian and Oceanian regions. *Mem Queensl. Mus.* **1989**, *26*, 521.
118. Pérez-Staples, D.; Prabhu, V.; Taylor, P.W. Post-teneral protein feeding enhances sexual performance of Queensland fruit flies. *Physiol. Entomol.* **2007**, *32*, 225–232. [[CrossRef](#)]
119. Kováts, E. Gas-chromatographische Charakterisierung organischer Verbindungen. Teil 1: Retentionsindices aliphatischer Halogenide, Alkohole, Aldehyde und Ketone. *Helvetica Chim. Acta* **1958**, *41*, 1915–1932. [[CrossRef](#)]
120. McCarthy, E.D.; Han, J.; Calvin, M. Hydrogen atom transfer in mass spectrometric fragmentation patterns of saturated aliphatic hydrocarbons. *Anal. Chem.* **1968**, *40*, 1475–1480. [[CrossRef](#)]

121. Nelson, D.R.; Sukkestad, D.R. Normal and branched aliphatic hydrocarbons from the eggs of the tobacco hornworm. *Biochem.* **1970**, *9*, 4601–4611. [[CrossRef](#)]
122. Nelson, D.R.; Sukkestad, D.R.; Zaylskie, R.G. Mass spectra of methyl-branched hydrocarbons from eggs of the tobacco hornworm. *J. Lipid Res.* **1972**, *13*, 413–421.

Sample Availability: Samples of the compounds are not available from the authors.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).