

Congenital tricuspid valve dysplasia in a septuagenarian

Lynn Khor ^{1*}, Michael Vallely ², and Martin Brown ²

¹Department of Cardiology, Macquarie University Hospital, 3 Technology Place, Macquarie University, NSW, 2109, Australia; and ²Department of Cardiology, Faculty of Medicine and Health Sciences, Macquarie University, NSW, Australia

Received 26 January 2020; first decision 31 March 2020; accepted 13 July 2020; online publish-ahead-of-print 10 September 2020

Primary isolated tricuspid regurgitation (TR) can be due to congenital or acquired tricuspid valve (TV) pathology. Anatomic studies suggest great variability in number and morphology of TV cusps with only 62% having three cusps.¹ We describe the case of a 77-year-old woman presenting with her first episode of right heart failure secondary to severe TR. Transoesophageal echocardiography revealed a congenital bileaflet tricuspid valve. The patient underwent surgical TV replacement, visually confirming fusion of the anterior and posterior leaflets of her TV with no other congenital abnormalities.

A 77-year-old woman presented with exertional dyspnoea and symptoms of right heart failure (New York Heart Association III–IV). Her comorbidities include chronic atrial fibrillation, obstructive sleep

apnoea, non-obstructive coronary artery disease, hypertension, and asthma. She was a lifelong non-smoker. Blood results including liver function tests were normal.

Transthoracic echocardiogram showed an apparently normal tricuspid valve (TV) with moderate-to-severe regurgitation, mildly dilated right ventricle (RV) with normal systolic function, mildly dilated right atrium (RA), borderline pulmonary pressure [right ventricular systolic pressure (RVSP 35 mmHg)], and normal left ventricular size and function.

Coronary angiogram showed minor coronary disease with normal left ventricular (LV) function and LV end-diastolic pressure (10 mmHg). Right heart catheterization showed normal pulmonary and




Figure 1 Three-dimensional transoesophageal echocardiography (TOE) showing atrial view of the tricuspid valve (TV) with two leaflets. AP, fused anterior and posterior leaflets; S, septal leaflet.




Figure 2 Intraoperative photograph of tricuspid valve with fusion of the anterior and posterior leaflets. AP, fused anterior and posterior leaflets; S, septal leaflet.

*Corresponding author. Tel: 612 4734 1000, Email: lynn.j.khor@gmail.com

Handling Editor: Hatem Soliman Aboumarie

Peer-reviewer: Christoph Sinning

© The Author(s) 2020. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

RA pressures (6 mmHg), although large tricuspid regurgitation (TR) waves were evident, and no left to right shunt. Her cardiac output was reduced at 3.53 L/min (cardiac index 2.11 L/min/m²) by thermodilution.

Transoesophageal echocardiogram showed bileaflet (fusion of anterior and posterior leaflets) TV, with no prolapse or flail leaflets (Figure 1). There was moderate-to-severe TR with moderately dilated RV and preserved systolic function.

As no other cardiac or respiratory cause for her dyspnoea was found, she proceeded to surgical TV replacement with a 29 mm Edwards Magna Ease prosthesis. Intraoperative findings confirmed the presence of a bileaflet TV (Figure 2). No other congenital abnormalities were identified.

This case highlights the importance of congenital TV abnormalities as a differential for primary isolated TR, resulting in RV and RA dilatation, with right heart failure, regardless of the age at presentation. We hypothesize that congenital TV pathology can present later in life

due to the lower right-sided pressures. While functional TR remains the most prevalent aetiology, primary isolated TR (both congenital and acquired) still accounts for 10% of TR.² Tricuspid valve surgery remains the definitive therapy for right heart failure from primary isolated TV disease.

Consent: The author/s confirm that written consent for submission and publication of this case report including image(s) and associated text has been obtained from the patient in line with COPE guidelines.

Conflict of interest: none declared.

References

1. Gerola LR, Wafae N, Vieira MC, Juliano Y, Smith RBW, Prates J. Anatomic study of the tricuspid valve in children. *Surg Radiol Anat* 2001;**23**:149–154.
2. Taramasso M, Pozzoli A, Guidotti A, Nietlispach F, Inderbitzin DT, Benussi S et al. Percutaneous tricuspid valve therapies: the new frontier. *Eur Heart J* 2017;**38**: 639–647.