

RESEARCH ARTICLE

Open Access

Control of the olive fruit fly using genetics-enhanced sterile insect technique

Thomas Ant^{1,2}, Martha Koukidou¹, Polychronis Rempoulakis^{1,3}, Hong-Fei Gong¹, Aris Economopoulos³, John Vontas³ and Luke Alphey^{1,2*}

Abstract

Background: The olive fruit fly, *Bactrocera oleae*, is the major arthropod pest of commercial olive production, causing extensive damage to olive crops worldwide. Current control techniques rely on spraying of chemical insecticides. The sterile insect technique (SIT) presents an alternative, environmentally friendly and species-specific method of population control. Although SIT has been very successful against other tephritid pests, previous SIT trials on olive fly have produced disappointing results. Key problems included altered diurnal mating rhythms of the laboratory-reared insects, resulting in asynchronous mating activity between the wild and released sterile populations, and low competitiveness of the radiation-sterilised mass-reared flies. Consequently, the production of competitive, male-only release cohorts is considered an essential prerequisite for successful olive fly SIT.

Results: We developed a set of conditional female-lethal strains of olive fly (named Release of Insects carrying a Dominant Lethal; RIDL[®]), providing highly penetrant female-specific lethality, dominant fluorescent marking, and genetic sterility. We found that males of the lead strain, OX3097D-Bol, 1) are strongly sexually competitive with wild olive flies, 2) display synchronous mating activity with wild females, and 3) induce appropriate refractoriness to wild female re-mating. Furthermore, we showed, through a large proof-of-principle experiment, that weekly releases of OX3097D-Bol males into stable populations of caged wild-type olive fly could cause rapid population collapse and eventual eradication.

Conclusions: The observed mating characteristics strongly suggest that an approach based on the release of OX3097D-Bol males will overcome the key difficulties encountered in previous olive fly SIT attempts. Although field confirmation is required, the proof-of-principle suppression and elimination of caged wild-type olive fly populations through OX3097D-Bol male releases provides evidence for the female-specific RIDL approach as a viable method of olive fly control. We conclude that the promising characteristics of OX3097D-Bol may finally enable effective SIT-based control of the olive fly.

Keywords: olive fly, *Bactrocera oleae*, sterile insect technique, SIT, release of insects carrying a dominant lethal, RIDL, autocidal control, insect transgenics

Background

The olive fly, *Bactrocera oleae* (Rossi) (Diptera: Tephritidae), is the major insect pest of olives. Females typically lay a single egg per olive [1], injected into the fruit through the female's ovipositor. The developing larva tunnels through the olive, feeding on the fleshy mesocarp. Heavy olive fly infestation can reduce the quality and therefore the value of the olive oil by up to 80%, and cause the

rejection of entire harvests of table olives [2]. Control currently relies overwhelmingly on the use of chemical insecticides, and because of the high economic and environmental costs of chemical control, together with the appearance of insecticide-resistant populations [3], there is an urgent need for improved control methods.

The sterile insect technique (SIT) is an environmentally friendly and species-specific method of pest control based on the release of large numbers of sterilised insects [4]. Competition for mating between wild and sterile males results in a decrease in the number of fertile matings, and

* Correspondence: luke.alphey@oxitec.com

¹Oxitec Limited, 71 Milton Park, Oxford OX14 4RX, UK

Full list of author information is available at the end of the article

a decline in the overall population size. SIT has been successfully implemented against various pest insect species including several Tephritidae. However, despite decades of research aimed at developing an olive fly SIT programme using radiation-sterilised flies [5], the consistently poor results led to eventual abandonment of the trials. Key issues included low quality of the radiation-sterilised mass-reared flies, economical production of sufficient numbers of sterilised flies, and assortative mating of released and wild populations because of different preferred mating times [5-7]. Laboratory-reared wild-type flies were found to mate several hours earlier than wild flies [7] (presumably due to differential selective pressures in the artificial laboratory-rearing environment). The proposed solution was male-only release [6]. Male-only release using a genetic sexing strain has also been shown to give a threefold to fivefold improvement in the performance of released radiation-sterilised *Ceratitidis capitata* (Medfly) males used for SIT [8].

The RIDL[®] (Release of Insects carrying a Dominant Lethal) system is a transgene-based derivative of SIT [9-14], one version of which involves the mass release of insects carrying a female-specific lethal transgene (fsRIDL) [15-17]. Encouraged by work advancing olive fly mass rearing by the International Atomic Energy Agency (IAEA) and others, we set out to develop fsRIDL strains of olive fly to overcome the remaining historic limitations of olive fly SIT. We therefore transformed the olive fly with construct OX3097, an fsRIDL system that had previously provided suitable fsRIDL strains for Medfly, incorporating genetic sexing, genetic sterilisation, and a heritable fluorescent marker [15].

We report the development of the first transgenic control strains for the olive fly. The lead strain, OX3097D-Bol, provides highly penetrant, dominant, female-specific lethality when reared in the absence of a transgene repressor. Using a series of behavioral studies, we found that OX3097D-Bol males show fully synchronous and strongly competitive mating behavior with wild olive flies, and display no reduction in their ability to induce refractoriness to wild female re-mating. This is a more stringent test than using laboratory wild-type flies, which may have co-adapted to laboratory rearing conditions. Furthermore, we describe results from experiments showing the ability of periodic releases of OX3097D-Bol males to suppress large caged wild-type olive fly populations.

Results

Olive fly transformation with OX3097 and phenotypic analysis


Six transgenic lines were generated carrying the OX3097 transgene (Figure 1A). Of the five lines with single insertions, two showed female-specific lethality that was fully penetrant and efficiently repressed by dietary tetracycline


(Figure 1C, D). Of these, line OX3097D-Bol (Figure 1E), which also showed the brightest expression of the DsRed2 fluorescent marker, was selected for further analysis. Consistent with previous observations of the OX3097 construct in Medfly [15] and Mexfly (Stainton *et al.*, unpublished data), female lethality occurred at the early pupal stage. Analysis of transgene-derived transcripts indicated that the alternative splicing pattern of the *Cctra* intron in OX3097D-Bol olive fly is equivalent to that in its native context [15,18,19] (Figure 1B). The OX3097D-Bol strain was selected for further development, and made homozygous for the transgene.

Mating tests with OX3097D-Bol and wild olive flies

Caged mating competitiveness tests challenging homozygous OX3097D-Bol males to compete with wild males for copulations with wild females were performed based on established guidelines [20], using wild olive flies from infested olives collected in Crete. In total, 15 experiments were performed, each with 50 OX3097D-Bol males, 50 wild males, and 50 wild females, thus more than 400 total couples were assessed. Wild males outperformed OX3097D-Bol males, gaining an average of 56% of total mates ($P = 0.01$, degrees of freedom (d.f.) = 1, likelihood ratio test for goodness of fit, $n = 406$). Nonetheless, this near-equal outcome far exceeds established thresholds for SIT [20]. Copulation initiation times were also recorded (Figure 2A). No significant difference was seen in mating initiation times for each type of male ($P = 0.44$, d.f. = 1, circular statistics F -test), suggesting synchronicity in mating activity between OX3097D-Bol males and wild flies.

Olive fly females typically only mate once [21]; males transfer more sperm than the female needs to fertilise all her eggs, and mating induces long-lasting refractoriness to re-mating, although if the females are held in close proximity to males some re-mating occurs. For Medfly, radiation sterilisation not only reduces the ability of male flies to compete for mates but also increases the likelihood that mated females will subsequently re-mate; that is, radiation reduces the ability of the males to induce refractoriness to re-mating in females [22]. Furthermore, the second mate taken is also preferentially wild rather than sterile. These effects of reduced male mating success, increased female re-mating, and preferential mating to wild (fertile) males on re-mating each reduce the efficiency of SIT. We therefore investigated the effect of first-male genotype on wild female re-mating propensity and second-mate genotype. Wild females who had initially mated either an OX3097D-Bol male ($n = 188$) or a wild male ($n = 296$) were caged with and allowed to mate with equal proportions of OX3097D-Bol and wild males over 15 days. No significant differences in re-mating rate ($P = 0.7$, d.f. = 1, χ^2 test), or second-mate genotype were seen ($P = 0.38$, d.f. = 1, χ^2 test) (Figure 2B).





Elimination of caged olive fly populations

We tested the ability of periodic release of OX3097D-Bol males to suppress target populations of wild-type olive fly, as published previously [17]. Wild-type populations totaling approximately 400 male and female olive flies were established in each of four large (8 m³) cages in a greenhouse at the University of Crete. At week 13 from first establishment, approximately 1,600 OX3097D-Bol male pupae were introduced per week. Population density was monitored by measuring egg production (Figure 3A) and female mortality (Figure 3B). To track mating outcomes, pupae taken from the cage were scored for fluorescence (Figure 3C). Fluorescent pupae were detected at three weeks post-first release (PR) and the proportion increased rapidly thereafter to 100% by week 10 PR. Death of fluorescent (transgenic) female pupae as a result of expression of the lethal phenotype would reduce the number of egg-laying adult females; based on the fluorescence data, no wild-type females emerged (eclosed) in the treatment cages from week 10 PR. Egg production was lower in the treatment than in the control cages in week 6 PR, and declined rapidly thereafter. The numbers of dead females recovered from treatment cages started to decrease from around week 6 to 7 PR, presumably due to a decline in overall female numbers. Defining extinction as 2 weeks of zero egg production [17], both treatment cage populations were extinct by week 12 PR.

Discussion

An effective non-chemical approach to olive fly management is highly desirable, and SIT could provide this if

specific technical issues can be overcome. Considerable progress has been made by others in respect of olive fly mass-rearing and release, and we set out to overcome the key remaining issues through genetics.

The use of genetic technology in food production has been controversial in some countries, notably within the European Union (EU). It is possible that the use of genetically modified arthropods to protect the crop in a sustainable way, which leaves little or no residue in the food product (here olives or olive oil) might be more acceptable in the EU than food products that are produced directly from genetically modified organisms. For comparison, the Standing Committee on the Food Chain and Animal Health stated that foods and feeds produced by fermentation with genetically modified microorganisms that are not present in the final product are "excluded from the scope of regulations" [23]. Additionally, it is recognized in EU legislation that the presence of technically unavoidable traces of genetically modified organisms in food and feed products should not trigger labeling requirements (Regulation 1830/2003/EC), as long as appropriate steps to avoid the presence of the materials have been taken. This is also analogous to the situation regarding the presence of wild-type insect parts as contaminants in food products [24].

Using modern genetic engineering techniques, we have developed OX3097D-Bol, an fsRIDL olive fly strain that incorporates engineered phenotypes, directly addressing problems encountered in previous unsuccessful olive fly SIT trials [5]. OX3097D-Bol is able to provide 100%




Figure 3 Population elimination by periodic release of OX3097D-Bol males. (A) The average daily egg production for each cage. Weeks 1 to 12 was the population stabilization period with 250 pupae added in the first week, and 200 pupae added to each cage per week thereafter. From week 13, 1,600 OX3097D-Bol pupae were added weekly into cages A and B. After week 13, weekly pupal return to the treatment cages was made proportional to the weekly egg production in the cage relative to the control cages. From 5 weeks after initiation of RIDL introductions, egg production in each treatment cage was consistently lower than in either control cage; the difference increased until eventual extinction of the wild-type population in both treatment cages by week 24 (12 weeks after the first RIDL release). Extinction was defined as 2 weeks of zero egg production. Egg numbers in control cages remained relatively stable. (B) Dead flies were removed from the cages weekly, and the numbers of dead females are shown. From 7 weeks after the initiation of RIDL release, increasingly fewer such females were recovered from the treatment cages than from the control cages. (C) Frequency of DsRed2 in treatment cages. Larvae selected for return were screened for presence of DsRed2 marker by fluorescence microscopy before being returned to the treatment cage (see Methods). The proportion of returning pupae carrying the OX3097D-Bol transgene reached 100% in both treatment cages by week 23 (10 weeks post-RIDL release). Olive fly females typically mate only once [21] (Figure 2B). Females start to lay eggs approximately 2 days after mating, and lay most of their eggs within the next 10 days. Egg to pupa development time was approximately 12 days. These pupae therefore indicate female mating choice of approximately 3 weeks before each measurement.

male-only release generations when reared in the absence of a dietary transgene repressor. Genetic sexing encourages random mating between the wild and the

released sterile populations, providing established performance benefits [8]. Female lethality renders redundant the requirement of performance-reducing and costly

sterilisation using ionising radiation. Furthermore, the dominant fluorescent marker has the potential to improve field monitoring, and the requirement of tetracycline for female development mitigates risks of accidentally released transgenic populations establishing in the wild. Introgression of susceptibility alleles through the female line provides a resistance management option for other control methods used with fsRIDL in an integrated pest-management programme [25-27].

Conclusions

The good mating characteristics of OX3097D-Bol males seen in this study, including high sexual competitiveness, photo-period compatibility, and efficient induction of re-mating refractoriness in wild females, suggest a strong potential for application in a SIT control programme. Although the results presented here need field confirmation, the proof-of-principle suppression and elimination of caged wild-type olive fly populations through OX3097D-Bol male releases provides evidence for the fsRIDL approach as a viable method for olive fly control. Independently, significant advances have been achieved in olive fly mass-rearing [28,29] and release [30] techniques. We therefore propose that the development of OX3097D-Bol will help form the basis of more effective and more sustainable control of this ancient and destructive olive pest.

Methods

Olive fly strains, rearing, and transformation

Olive fly rearing was carried out using standard methods. Approximately 4,500 pre-blastoderm stage olive fly embryos of the Democritus laboratory (Greece) *B. oleae* strain were micro-injected with the OX3097 plasmid and *piggyBac* mRNA, as described by Koukidou *et al.* [31]. This resulted in 138 surviving G₀ adults. After back-crossing pools of five male survivors with ten wild-type females, and five female survivors with five wild-type males, six OX3097 lines were isolated (transformation efficiency ~4%). Transgenics were outcrossed for five generations to the Argov wild-type strain (Israel), and these outcrossed Argov derivatives were used for all experiments. The Argov strain was derived from field collections of wild male olive fly in Israel, outcrossed to Democritus females. To develop a homozygous strain, a pool of homozygous and heterozygous OX3097D-Bol adults was generated by crossing OX3097D-Bol heterozygotes. DNA from these parents was analysed by PCR using primers (5'-CCTGCGTTTGGAGATGACGAAATC-3' and 5'-CTTACATATAGAGCAGTGCGCTCATG-3') that anneal to genomic sites flanking the insertion site producing a WT (no insertion) amplicon were discarded. Homozygotes were thereby identified, and a homozygous line was developed from 15 female and

13 male founder flies. The key phenotypic properties identified in OX3097D-Bol heterozygotes (Figure 1, 2) were also found in the homozygous line.

Reverse transcriptase PCR

Male and female tTAV [14,34] transcripts were amplified by reverse transcriptase (RT)-PCR using cDNA synthesized using a commercial kit (Superscript II; Life Technologies, Grand Island, NY, USA) on RNA extracts prepared using TRI reagent (Life Technologies). The major RT-PCR products were sequenced (GATC Biotech, Konstanz, Germany) and analysed using VectorNTi (Life Technologies).

Mate competition and re-mating tests

Mating competitiveness tests were carried out in accordance with the United Nations Food and Agriculture Organization (FAO)/IAEA/United States Department of Agriculture (USDA) guidelines [20] in semi-natural caged conditions (cages were 1.25 m high with a base of 0.25 m² and contained a large olive branch) in greenhouse facilities at the University of Crete under natural light. Adult male OX3097D-Bol flies were obtained from larvae reared in the absence of tetracycline ("off-tet") at low density (1 larva/0.8 g larval medium). Wild pupae were recovered from infested olives gathered from olive orchards near to the University of Crete. Each mating test used 50 OX3097D-Bol males, 50 wild males, and 50 wild females. Mated males were scored for the presence of the DsRed2 fluorescent marker by epifluorescence microscopy [32,33] (Clontech Laboratories, Inc, Mountain View, CA, USA). OX3097D-Bol mating competition tests were performed in 15 replicates with more than 400 couples assessed in total. Each experiment yielded a mating propensity of greater than 0.2 [20]. Mating cages for the first step of re-mating tests contained either wild females and OX3097D-Bol males or wild females and wild males. Mating couples were removed from cages. Mated females were grouped in accordance with the genotype of the first mate (OX3097D-Bol, n = 188 or wild-type, n = 296) and were transferred the following day to new cages with sufficient fresh wild and OX3097D-Bol males to give a 1:1:1 ratio of mated females, wild males, and OX3097D-Bol males. Cages were checked daily for re-mating events over the following 15 days. Re-mating couples were removed, and the male genotype assessed by epifluorescence microscopy.

Caged suppression of stable wild-type populations

The protocol for the caged suppression experiment was based on that of Wise de Valdez *et al.* [17]. Stable populations of wild-type (Argov) olive flies were established in four field cages (each 8 m³ and containing an olive tree 1.5 m tall; all the cages were contained within a single

large glasshouse) over a 12-week period by introducing a fixed number of pupae to each cage weekly (250 in week 1, and 200 from weeks 2 to 12). To assess egg production rates and to sustain caged olive fly populations, four cerasin-wax cones (combined surface area approximately 550 cm²) were added to each cage daily for female oviposition. Pupal additions for the first 4 weeks during population establishment originated from a wild-type stock colony; thereafter, experimental cages were self-sufficient. Eggs were collected from the cages and counted daily. On week 12, the experimental cages were randomly divided into two RIDL-treatment and two no-treatment (control) cages. From week 13 onwards, RIDL-treatment cages received weekly additions of 1,600 OX3097D-Bol male pupae reared off-tet (an initial recruitment rate ratio of approximately 16 OX3097D-Bol males to 1 wild-type male). Once the OX3097D-Bol introductions began, pupal introductions to the RIDL-treatment cages were proportional to the cage's respective rate of egg production, with the control cages providing a coefficient of weekly egg production to pupal return. Numbers of females in a cage were monitored by collecting and counting dead females. The ratio of RIDL heterozygous to wild-type pupae returning to RIDL-treatment cages was monitored by fluorescence microscopy (OX3097D-Bol heterozygous females reared off-tet pupate but fail to eclose). Larvae were screened only after the returning population was separated, to remove the possibility of bias in selecting individual flies for reintroduction to the cage populations.

Statistical analysis

Comparison of mating competitiveness between OX3097D-Bol and wild males was performed using a likelihood ratio test for goodness of fit with numbers of successfully mated males from both genotypes pooled across experiments, and compared against a theoretical 1:1 genotype ratio expected in random mating. Differences in copulation initiation time between the two possible mating combinations (OX3097D-Bol/wild female and wild male/wild female) were analysed using a circular statistics *F*-test with the statistical software Oriana for Windows (Kovach Computing Services, Anglesey, UK). The Pearson's χ^2 test was used to compare numbers of re-mated to non-re-mated females for both initial mating combinations. The same test was used to compare initial and second-mate choice. Unless otherwise stated, all statistical tests were performed using SPSS for Windows (version 10.0; SPSS Inc., Chicago IL USA), with significance set at $P < 0.05$.

Abbreviations

CcTra intron: *Ceratitis capitata* transformer intron; fsRIDL: female-specific Release of Insects carrying a Dominant Lethal; PR: post-first-release; RIDL: Release of Insects carrying a Dominant Lethal; SIT: Sterile Insect Technique.

Acknowledgements

We thank the Agricultural Research Organisation, Israel (ARO) and Bio-Fly Ltd for the Argov wild-type strain. We thank Zoe Lokhon, Chrysargyris Antonios, Sian Morgan, Camilla Beech, and the Oxitec molecular biology team for technical assistance. TA was supported by an Industrial Case DPhil studentship from the UK Biotechnology and Biological Sciences Research Council.

Author details

¹Oxitec Limited, 71 Milton Park, Oxford OX14 4RX, UK. ²Department of Zoology, University of Oxford, South Parks Road, Oxford OX1 3PS, UK. ³Faculty of Biotechnology and Applied Biology, Department of Biology, University of Crete, Heraklion, Crete, Greece.

Authors' contributions

TA and MK created the transgenic olive fly lines. TA conducted the molecular analysis. TA, MK, PR, carried out mating tests and suppression trial. HG, JV and AE provided advice, LA conceived the project, and MK supervised the project. TA and LA wrote the paper. All authors discussed results and commented on the manuscript.

Competing interests

TA, MK, PK, HFG and LA are employees or students of Oxitec Ltd and have employment, stipend and/or equity interest in Oxitec. Oxitec and the University of Oxford have intellectual property related to the subject matter of this paper. All other authors declare no interest.

Received: 9 March 2012 Accepted: 19 June 2012

Published: 19 June 2012

References

1. Burrack HJ, Fornell AM, Connell JH, O'Connell NV, Phillips PA, Vossen PM, Zalom FG: **Intraspecific larval competition in the olive fruit fly (Diptera: tephritidae).** *Environ Entomol* 2009, **38**:1400-1410.
2. Daane KM, Johnson MW: **Olive fruit fly: managing an ancient pest in modern times.** *Annu Rev Entomol* 2010, **55**:151-169.
3. Vontas J, Hernandez-Crespo P, Margaritopoulos JT, Ortego F, Feng H, Mathiopoulos KD, Hsu J: **Insecticide resistance in Tephritid flies.** *Pest Biochem and Physiol* 2011, **100**:199.
4. Dyck VA, Hendrichs J, Robinson AS: *Sterile Insect Technique: Principles and practice in area-wide integrated pest management* Dordrecht, Netherlands: Springer; 2005.
5. Economopoulos AP, Avtzis N, Zervas G, Tsitsipis J, Haniotakis G, Tsiropoulos G, Manoukas A: **Control of the olive fly, *Dacus oleae* (Gmelin), by the combined effects of insecticides and release of gamma sterilised insects.** *J Appl Entomol* 1977, **83**:201-215.
6. Economopoulos AP: **The olive fly, *Bactrocera (Dacus) oleae* (Gmelin) (Diptera:Tephritidae): its importance and control; previous SIT research and pilot testing.** Austria: International Atomic Energy Agency; 2001, 44.
7. Zervas GA, Economopoulos AP: **Mating frequency in caged populations of wild and artificially reared (normal or gamma-sterilized) olive fruit flies, *Dacus oleae* (Gmelin) (Diptera:Tephritidae).** *Environ Entomol* 1982, **11**:17-20.
8. Rendon P, McInnis D, Lance D, Stewart J: **Medfly (Diptera: Tephritidae) genetic sexing: large-scale field comparison of males-only and bisexual sterile fly releases in Guatemala.** *J Econ Entomol* 2004, **97**:1547-1553.
9. Alphey L, Nimmo D, O'Connell S, Alphey N: **Transgenesis and the management of vector-borne disease.** Edited by: Aksoy S. Berlin: Springer-Verlag; 2008:627:93-103.
10. Alphey L, Andreasen M: **Dominant lethality and insect population control.** *Mol Biochem Parasitol* 2002, **121**(2):173-178.
11. Alphey L: **Re-engineering the sterile insect technique.** *Insect Biochem Mol Biol* 2002, **32**(10):1243-1247.
12. Morrison NI, Franz G, Koukidou M, Miller TA, Saccone G, Alphey LS, Beech CJ, Nagaraju J, Simmons GS, Polito LC: **Genetic improvements to the sterile insect technique for agricultural pests.** *Asia Pac J Mol Bio* 2010, **18**:275-295.
13. Thomas DD, Donnelly CA, Wood RJ, Alphey LS: **Insect population control using a dominant, repressible, lethal genetic system.** *Science* 2000, **287**(5462):2474-2476.

14. Gong P, Epton MJ, Fu G, Scaife S, Hiscox A, Condon KC, Condon GC, Morrison NI, Kelly DW, Dafa'alla T, Coleman PG, Alphey L: **A dominant lethal genetic system for autocidal control of the Mediterranean fruit fly.** *Nat Biotechnol* 2005, **23**(4):453-456.
15. Fu G, Condon KC, Epton MJ, Gong P, Jin L, Condon GC, Morrison NI, Dafa'alla TH, Alphey L: **Female-specific insect lethality engineered using alternative splicing.** *Nat Biotechnol* 2007, **25**(3):353-357.
16. Fu G, Lees RS, Nimmo D, Aw D, Jin L, Gray P, Berendonk TU, White-Cooper H, Scaife S, Kim Phuc H, Marinotti O, Jasinskiene N, James AA, Alphey L: **Female-specific flightless phenotype for mosquito control.** *Proc Natl Acad Sci USA* 2010, **107**(12):4550-4554.
17. Wise de Valdez MR, Nimmo D, Betz J, Gong HF, James AA, Alphey L, Black WC4th: **Genetic elimination of dengue vector mosquitoes.** *Proc Natl Acad Sci USA* 2011, **108**(12):4772-4775.
18. Pane A, Salvemini M, Delli Bovi P, Polito C, Saccone G: **The transformer gene in *Ceratitis capitata* provides a genetic basis for selecting and remembering the sexual fate.** *Development* 2002, **129**(15):3715-3725.
19. Dafa'alla T, Fu G, Alphey L: **Use of a regulatory mechanism of sex determination in pest insect control.** *J Genet* 2010, **89**(3):301-305.
20. FAO/IAEA/USDA: *Product quality control and shipping procedures for sterile mass-reared tephritid fruit flies* Vienna, Austria: International Atomic Energy Agency; 2003.
21. Tzanakakis KE, Tsitsipis JA, Economopoulos AP: **Frequency of mating in females of olive fruit fly under laboratory conditions.** *J Econ Entomol* 1968, **61**:1309-1312.
22. Kraaijeveld K, Chapman T: **Effects of male sterility on female remating in the Mediterranean fruit fly, *Ceratitis capitata*.** *Proc Biol Sci* 2004, **271**:209-11.
23. European Union: **Report from the Commission to the Council and the European Parliament on the implementation of Regulation (EC) No 1829/2003 of the European Parliament and of the Council on genetically modified food and feed.** [http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:52006DC0626:EN:HTML].
24. European Union: **Commission Implementing Regulation (EU) No 543/2011 of 7 June 2011 laying down detailed rules for the application of Council Regulation (EC) No 1234/2007 in respect of the fruit and vegetables and processed fruit and vegetables sectors.** *Official Journal of the European Union* 2011, 543/2011.
25. Alphey N, Bonsall MB, Alphey L: **Modelling resistance to genetic control of insects.** *J Theor Biol* 2011, **270**(1):42-55.
26. Alphey N, Bonsall MB, Alphey L: **Combining pest control and resistance management: synergy of engineered insects with Bt crops.** *J Econ Entomol* 2009, **102**(2):717-732.
27. Alphey N, Coleman PG, Donnelly CA, Alphey L: **Managing insecticide resistance by mass release of engineered insects.** *J Econ Entomol* 2007, **100**(5):1642-1649.
28. Dimou I, Rempoulakis P, Economopoulos AP: **Olive fruit fly [*Bactrocera oleae* (Rossi) (Diptera:Tephritidae)] adult rearing diet without antibiotic.** *J Appl Entomol* 2009, **134**:72-79.
29. Estes AM, Nestel D, Belcari A, Jessup A, Rempoulakis P, Economopoulos AP: **A basis for the renewal of sterile insect technique for the olive fly, *Bactrocera oleae* (Rossi).** *J Appl Entomol* 2011, **136**:1-16.
30. Rempoulakis P, Nestel D: **Dispersal ability of marked, irradiated olive fruit flies [*Bactrocera oleae* (Rossi) (Diptera: Tephritidae)] in arid regions.** *J Appl Entomol* 2012, **136**:171-180.
31. Koukidou M, Klinakis A, Reboulakis C, Zagoraiou L, Tavernarakis N, Livadaras I, Economopoulos A, Savakis C: **Germ line transformation of the olive fly *Bactrocera oleae* using a versatile transgenesis marker.** *Insect Mol Biol* 2006, **15**(1):95-103.
32. Lukyanov KA, Fradkov AF, Gurskaya NG, Matz MV, Labas YA, Savitsky AP, Markelov ML, Zaraisky AG, Zhao X, Fang Y, Tan W, Lukyanov SA: **Natural animal coloration can be determined by a nonfluorescent green fluorescent protein homolog.** *J Biol Chem* 2000, **275**(34):25879-25882.
33. Matz MV, Fradkov AF, Labas YA, Savitsky AP, Zaraisky AG, Markelov ML, Lukyanov SA: **Fluorescent proteins from nonbioluminescent Anthozoa species.** *Nat Biotechnol* 1999, **17**(10):969-973.
34. Phuc HK, Andreassen MH, Burton RS, Vass C, Epton MJ, Pape G, Fu G, Condon KC, Scaife S, Donnelly CA, Coleman PG, White-Cooper H, Alphey L: **Late-acting dominant lethal genetic systems and mosquito control.** *BMC Biol* 2007, **5**:11.

doi:10.1186/1741-7007-10-51

Cite this article as: Ant et al.: Control of the olive fruit fly using genetics-enhanced sterile insect technique. *BMC Biology* 2012 **10**:51.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



[Search](#) [Workspace](#) [Ulrich's Update](#) [Admin](#)

Enter a Title, ISSN, or search term to find journals or other periodicals:

[▶ Advanced Search](#)

[Search Results](#)

JCR[®] Web [ingentaconnect™](#)
 Search MultiSearch: [ISSN Search](#) | [Title Search](#)

BMC Biology

Title Details



Lists

[Marked Titles](#) (0)

Search History

[1741-7007](#) - (1)

[Save to List](#)
[Email](#)
[Download](#)
[Print](#)
[Corrections](#)
[Expand All](#)
[Collapse All](#)

▼ Basic Description	
Title	BMC Biology
ISSN	1741-7007
Publisher	BioMed Central Ltd.
Country	United Kingdom
Status	Active
Start Year	2004
Frequency	Irregular
Language of Text	Text in: English
Refereed 	Yes
Abstracted / Indexed	Yes
Open Access 	Yes http://www.biomedcentral.com/bmcbiol/
Serial Type	Journal
Content Type	Academic / Scholarly
Format	Online
Explanation of Title Acronym	BioMed Central
Website	http://www.biomedcentral.com/bmcbiol/
Email	editorial@biomedcentral.com
Description	Features original research articles and methodology articles in any area of biology but with a focus on the biomedical sciences, including biochemistry, cell biology, chemical biology, computational biology, developmental biology, ecology, evolutionary biology, genetics, genomics, immunology, microbiology, molecular biology, neurobiology, pharmacology, physiology, plant biology and structural biology.
▶ Subject Classifications	
▶ Additional Title Details	
▶ Publisher & Ordering Details	
▶ Online Availability	
▶ Abstracting & Indexing	
▶ Other Availability	
▶ Demographics	
▶ Reviews	

[Save to List](#)
[Email](#)
[Download](#)
[Print](#)
[Corrections](#)
[Expand All](#)
[Collapse All](#)

[Contact Us](#) | [Privacy Policy](#) | [Terms and Conditions](#) | [Accessibility](#)

Ulrichsweb.com™, Copyright © 2016 ProQuest LLC. All Rights Reserved