



Mountain cloud forest and grown-shade coffee plantations: A comparison of tree biodiversity in central Veracruz, Mexico

Alfredo González-Zamora¹, Manuel Esperón-Rodríguez^{1,2}, Víctor L. Barradas^{1*}

¹ *Laboratorio de Ecofisiología Tropical. Instituto de Ecología. Universidad Nacional Autónoma de México. 04510. México, D.F. México*

² *Department of Biological Sciences. Faculty of Sciences and Engineering. Macquarie University. NSW 2109. Australia*

Abstract

Aim of study: The objective of this work is to compare tree diversity and richness among one grown-shade coffee plantation (CAE) and two sites of montane cloud forests, one preserved (MCF1) and other perturbed (MCF2). We also develop an analysis of the importance of coffee plantations as a refuge of tree species, holding a potential role for conservation.

Area of study: Our study area is the coffee region of Coatepec-Xico, in the state of Veracruz, Mexico.

Material and Methods: We compiled a list of all tree species in each site to determine tree diversity and floristic similarity (dis-similarity). We used different similarity indices and a cluster analysis to show relations among sites.

Main results: 2721 individuals from 154 species were registered in the montane cloud forests as a whole. In the grown-shade coffee plantation we registered 2947 individuals from 64 species. The most similar sites were the perturbed montane cloud forest and the grown-shade coffee plantation and the least similar were the preserved montane cloud forest and the grown-shade coffee plantation. The high biodiversity found in all sites and the differences in tree composition between the two montane cloud forests supports evidence of the ecosystems richness in the region.

Research highlights: Diversity differences among sites determine that the grown-shade coffee plantation is not substitute for montane cloud forest. CAE's are developed under similar environmental conditions than the MCF; therefore, coexistence and re-combination (replacement) of species make them particularly complementary. CAE's in Veracruz have a potential role as refuge for biodiversity.

Keywords: Agroforestry systems; floristic similarity; diversity; richness; biodiversity refuge.

Citation: González-Zamora, A., Esperón-Rodríguez, M., Barradas, V.L. (2016). Mountain cloud forest and grown-shade coffee plantations: A comparison of tree biodiversity in central Veracruz, Mexico. *Forest Systems*, Volume 25, Issue 1, e055. <http://dx.doi.org/10.5424/fs/2016251-07538>.

Received: 11 Feb 2015. **Accepted:** 09 Feb 2016

Copyright © 2016 INIA. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial (by-nc) Spain 3.0 Licence, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Funding: This research was supported by CONACyT (Mexican Council of Science and Technology) and SEMARNAT (Secretariat of Environment and Natural Resources) research grant SEMARNAT-2004-C01-332.

Conflicts of Interest: The authors have declared that no competing interests exist.

Correspondence should be addressed to Víctor L. Barradas: vlbarradas@ecologia.unam.mx

Introduction

In tropical regions, extensive conversion of forests and agricultural intensification are typically identified as the most prominent drivers of land-use change and biodiversity loss (Sala *et al.*, 2000; Wright 2005). The mitigation of tropical deforestation and biodiversity protection must address livelihoods and needs of local communities (Bhagwat *et al.*, 2008). In this sense, agroforestry is considered as a promising approach to reduce deforestation and improve rural livelihoods in the tropics (Current *et al.*, 1995; Ashley *et al.*, 2006). Agroforestry is a land-use management system where trees or shrubs develop around or among crops or pas-

tureland, providing economic, social and environmental benefits (McNeely, 1995; Bhagwat *et al.*, 2008).

Agroforestry systems are often very small in size and surrounded by open landscapes and resemble forest fragments. Species distinctiveness (presence of rare or endemic species) is frequently low, even though their species richness (total number of species) might be equal to, or higher than that of neighboring forests (O'Dea & Whittaker, 2007). Many agroforestry systems are important for protection of species and habitats outside protected areas, and agroforestry systems can be considered as refuges for biodiversity (Bhagwat *et al.*, 2008; Manson *et al.*, 2008; Nonato de Souza *et al.*, 2012). These systems conserve biodiversity in remnant

habitats and provide potential movement for species among these remnants (Bhagwat *et al.*, 2008). These systems also provide environmental services such as carbon stock and sequestration (Albrecht & Kandji, 2003; Dávalos Sotelo *et al.*, 2008; Thangata & Hildebrand, 2012), improvement of environmental quality (Tornquist *et al.*, 1999; Geissert & Ibáñez, 2008), water harvesting, reducing water runoff, and increased recharge of aquifers, reduction of floods and droughts, among other services (Mejía *et al.*, 2004).

Almost three quarters of the planet's surface and 67.3% of Mexico's surface are covered by ecosystems managed or modified by humans (Pimentel *et al.*, 1992; McNeely, 1995; Palacio-Prieto *et al.*, 2000; SAGARPA, 2007). Because of the dominance of these systems, their management changes can affect biodiversity conservation and ecosystem services (Tilman *et al.*, 2002; MEA, 2005). For the montane cloud forest (MCF), despite its high strategic value for sustainable development, the key role it plays in the hydrological cycle, and being considered as reservoir of endemic biodiversity (Toledo-Aceves *et al.*, 2011), in Mexico it is considered the most threatened terrestrial ecosystem because of land-use changes and the effects of global climate change. Currently, this ecosystem has been assigned as high priority for conservation and promotion of sustainable development (Aldrich *et al.*, 1997; CONABIO, 2010; Toledo-Aceves *et al.*, 2011; Calderon Aguilera *et al.*, 2012).

Montane cloud forest (MCF) in Veracruz occupies ca. 1243.65 km² (1.73% of the total area; Ortega & Castillo, 1996; Ellis & Martinez, 2010; Castillo-Campos *et al.*, 2011). In the central region of the state the MCF area was reduced gradually because of the expansion of the coffee cultivation. Since the late XIX century to 1960's, MCF's were replaced with coffee agro-ecosystems (Ruelas-Monjardín *et al.*, 2014), and the forest fragmentation was accelerated because of the demographic pressure and territorial expansion (Williams-Linera *et al.*, 2002), where the greatest impact on vegetation (transformation in species composition) was caused mostly by deforestation, fires, plantations establishment, and land-use conversion to pasture (Ellis & Martinez, 2010). As a result the development of several types of grown-shade coffee such as shade monoculture, simple polyculture, diverse or traditional polyculture, and rustic plantations in which the forest canopy is used as shade for coffee have taken place.

Currently, Veracruz is the second largest producer of coffee in Mexico with the 24.7% of the national coffee production, occupying an area of 1520 km², equivalent to 13.92% of total of vegetation present in the state (Olguín *et al.*, 2011). Coffee agro-ecosystems (CAE) are developed at the lowest elevation of the MCF under similar environmental and climate conditions; therefore, coexistence and recombination (re-

placement) of species make them particularly complementary (Castillo-Campos *et al.*, 2011). When coffee plantations are under shade, the system "CAE-MCF" maintains forest cover, although with less species diversity compared to the undisturbed MCF. However, because of its structure, species diversity, and environmental services provided, CAE's are of great importance for conservation (Ellis & Martinez, 2010; Olguín *et al.*, 2011; Toledo-Aceves *et al.*, 2011).

In this work, we compared tree diversity and richness among one grown-shade coffee plantations (CAE) and two sites of montane cloud forests (MCF), one preserved (MCF1) and other perturbed (MCF2), in the coffee region of Coatepec-Xico, Veracruz, Mexico. We also analyzed the importance of coffee plantations as a refuge of tree species, holding a potential role for conservation.

Material and methods

Study area and site selection

The coffee region of Coatepec-Xico is located in the central highlands of the state of Veracruz, Mexico (19° 29' 25''N, 97° 02' 30''W). In this region, the MCF is the dominant vegetation type. The area is located in the eastern slope of the Cofre de Perote, with altitudes from 1000 to 1350 m asl. Climate is temperate humid with an average annual temperature of 18 °C and annual precipitation between 1000 and 1500 mm (CONABIO, 2010; González-Espinosa *et al.*, 2012). Dominated soil types are yellowish soils derived from volcanic rocks (Gómez-Pompa, 1978). The coffee region is located near the city of Coatepec in central Veracruz. Coatepec is the largest coffee producer of the state, with 24.59% of the total cultivated area in Veracruz (Landeros-Sánchez *et al.*, 2011; Olguín *et al.*, 2011).

We selected three sites. The first site (MCF1) corresponded to a preserved forest located at La Cortadura (19°29' 29''N, 97° 01' 58''W). The second site (MCF2, 19°26' 29''N, 97° 00' 02''W) was a perturbed forest, finding vegetation disturbance by anthropogenic causes with presence of *Citrus* spp. and some species of primary succession such as *Senecio arbore-scens* and *Myrsine coriacea*. The third site, a coffee agro-ecosystem (CAE) was located near to La Orduña (19° 29' 17''N, 97° 55' 32''W; Figure 1).

Data collection

For each MCFs site we sampled an area of 1500 m² with two perpendicular and two parallel transects. For the CAE site we also sampled an area of 1500 m² divided in ten traditional coffee areas dedicated only to




Figure 1. Study area and the study sites: undisturbed montane cloud forest (MCF1), perturbed montane cloud forest (MCF2), and coffee agro-ecosystem (CAE) within the coffee region of Coatepec-Xico located in the central highlands in the state of Veracruz, Mexico.

the grown-shaded coffee cultivation. We used the method of sample collection established by Mostacedo & Fredericksen (2000) and CONAFOR (2011). In the field, we determined the most dynamic areas of change and the more complex vegetative structures to be sampled and made transects of 15 x 100 m using GIS. This method was applied for the three sites. Sampling transects ended when new species were not found, and only repeated species were counted.

We counted all tree individuals within the sites, categorizing them *in situ* into two size classes: *i*) woody species with diameter at breast height (DBH) < 10 cm, and *ii*) woody species with DBH > 10 cm. All individuals counted at each site were identified to species level following the nomenclature of the Flora of Veracruz (Sosa & Gomez-Pompa, 1994) and the classifications of Cronquist (1988). Specimens that could not be identified in the field were collected and subsequently identified using taxonomic keys in the laboratory of Tropical Ecophysiology of the Institute of Ecology, Universidad Nacional Autónoma de México (UNAM).

Data analysis

All individuals found and counted were taken into account for the diversity estimation. The compiled list of all tree species was used to calculate species richness, diversity indices, and floristic similarity (dissimilarity) for all study sites. Diversity was determined considering

the number of tree species per unit of sampled area. Species diversity of each site was determined using the indices of dominance and equity of Margalef (DMg) and Simpson (λ), equity of Menhinick (DMn), and Shannon-Wiener; this in order to obtain the diversity parameters of species and their quantification and representativeness (Mostacedo & Fredericksen, 2000; Moreno, 2001; Villarreal *et al.*, 2004). To compare the number of species shared among sites, we estimated the floristic similarity using the similarity/dissimilarity indices and the coefficients of Jaccard (IJ), Sørensen (IS), Morisita-Horn (IM-H), and the similarity coefficient of Sørensen for quantitative data (Iscuant). All data were entered with the established formulas of diversity indices to a database where calculations were performed to determine the diversity in *Excel version 14.5.2*.

For a visual representation of the potential relationships among sites and to determine whether the degree of environmental disturbance of each site allowed a specific grouping we plotted a cluster dendrogram. In this case we used as measure the distance of Manhattan and the Average method. All statistical analyses were conducted using the statistical environment software R (RCoreTeam, 2014).

Results

The most representative/abundant species were: *i*) MCF1: *Parathesis melanosticta* and *Hedyosmum mexicanum*, *ii*) MCF2: *Beilschmiedia mexicana*, *Clethra macrophylla* and *Carpinus tropicalis*, and *iii*) CAE:

Citrus spp. and *Inga vera* (Table 1). For MCF1 the 14 most abundant species accounted only 15.26 %; whereas for MCF2 and CAE the 14 most abundant species accounted 70 and 77.5% respectively (Table 1). For MCF2 we found several species that evidenced the perturbation degree: *Citrus* spp., *Heliocarpus donnell-smithii*, *Lippia myriocephala*, *Myrsine coriácea*, *Solanum schlechtendalianum* and *Trema micrantha*.

2721 individuals from 154 tree species were registered in MCF1 and MCF2; 116 species in MCF1 and 38 in MCF2. In CAE we registered 64 tree species with 2947 individuals (Table 1). The highest and lowest species diversity corresponded to MCF1 and MCF2 respectively; this was reflected in the Margalef (DMg) and the Menhinick (DMn) indices (Table 2). The Shannon-Wiener (H) index also indicated that MCF1 had the greatest diversity of species, and the Simpson dominance (λ) index showed that CAE was the least diverse site, whereas MCF1 was the most diverse (Table 2).

Regarding floristic similarity, we found that MCF1 and MCF2 shared 15 species, with *Alchornea latifolia*, *Psychotria* sp. and *C. macrophylla* as the most represented (higher number of individuals; Table 3). The greatest similarity was found between sites MCF2 and CAE, sharing 16 species, of which the most frequent species were *C. tropicalis* and *Citrus* spp. (Table 3). This similarity between MCF2 and CAE was also observed in the cluster analysis (Figure 2). The sites sharing less number of species (12) were MCF1 and CAE (Table 3). The Jaccard (IJ) and the Sørensen (IS) similarity coefficients, and the Morisita-Horn (IM-H) and the Sørensen (Iscuant) indices also confirmed that sites with less similarity were MCF1 and CAE, and those having the highest similarity were MCF2 and CAE. Values obtained to calculate these indices were higher for both qualitative and quantitative data (Table 4).

Discussion

We confirmed similarity/dissimilarity among sites using different qualitative and quantitative methods. Measuring species' relative abundance and similarity among sites allowed us to identify those species whose low representation make them more sensitive or vulnerable to environmental perturbations: *B. mexicana*, *Miconia glaberrima* and *I. punctata*. It is important to note that the most similar sites were MCF2 and CAE, but only with 16 species, i.e. the 20% of species present in MCF2 and CAE as a whole. Low similarity among sites might be due to the low number of common species between pairs of comparisons, finding a high percentage of species exclusive of each site, which contributes to biodiversity conservation at regional scale. These findings highlight the CAE importance for conservation and the high tree biodiversity in the region, especially considering that the sites are not far apart geographically (Figure 1).

Another important finding is the high dissimilarity between MCF1 and MCF2, where the low number of species in MCF2 shows the shocking biodiversity loss in perturbed areas. Although ecosystems such as MCF2, are subject to influences determined by other species (e.g. predators, competitors, invaders), and temporal and spatial variations of environmental conditions, such as nutrient availability, temperature and precipitation (Chapin *et al.*, 2000; Bellemare *et al.*, 2002), human activities and perturbations have a great impact on them. Human perturbations can decrease local diversity or richness, as it was seen in MCF2; however, for CAE this is not necessarily true. Human perturbations can widely change floristic composition of ecosystems, but agroforestry can help to mitigate



Figure 2. Cluster analysis showing the relationships among our study sites: undisturbed montane cloud forest (MCF1), perturbed montane cloud forest (MCF2), and coffee agro-ecosystem (CAE) within the coffee region of Coatepec-Xico located in the central highlands in the state of Veracruz, Mexico.

Table 1. Total number of individuals and percentage of the most abundant species of the three study sites from the coffee region of Coatepec-Xico, Veracruz, Mexico: *i*) undisturbed montane cloud forest (MCF1); *ii*) perturbed montane cloud forest (MCF2), and *iii*) coffee agro-ecosystem (CAE)

Species	Number of individuals	Percentage (%)
MCF1		
<i>Zanthoxylum melanostictum</i> Schltdl. & Cham	10	0.72
<i>Phyllonoma laticuspis</i> (Turcz.) Engl.	10	0.72
<i>Arachnothryx bourgaei</i> (Standl.) Borhid	10	0.72
<i>Oreopanax xalapensis</i> (Kunth) Decne. & Planch	12	0.86
<i>Clethra macrophylla</i> DC.	12	0.86
<i>Turpinia occidentalis</i> (Swartz) G. Don.	13	0.94
<i>Calyptranthes schlechtendaliana</i> O. Berg	13	0.94
<i>Miconia glaberrima</i> (Schltdl.) Naudin	16	1.15
<i>Alchornea latifolia</i> Sw.	17	1.22
<i>Piper xanthostachyum</i> C. DC.	17	1.22
<i>Psychotria</i> spp.	17	1.22
<i>Miconia chrysonera</i> Triana	18	1.30
<i>Hedyosmum mexicanum</i> Cordem.	22	1.59
<i>Parathesis melanosticta</i> (Schltdl.) Hemsl.	25	1.80
Number of individuals with the highest frequency	212	15.27
Total number of individuals (N)	1388	-
MCF2		
<i>Brunellia mexicana</i> Standl.	38	2.85
<i>Quercus xalapensis</i> Bonpl.	38	2.85
<i>Senecio arborescens</i> Steetz	39	2.93
<i>Quercus leiophylla</i> A. DC.	44	3.30
<i>Myrsine coriacea</i> (Sw.) R. Br. ex Roem. & Schult.	46	3.45
<i>Liquidambar styraciflua</i> L.	47	3.53
<i>Styrax glabrescens</i> Benth.	47	3.53
<i>Citrus</i> spp.	51	3.83
<i>Hampea integerrima</i> Schltdl.	51	3.83
<i>Quercus insignis</i> M. Martens & Galeotti	60	4.50
<i>Turpinia insignis</i> (Kunth) Tul.	79	5.93
<i>Beilschmiedia mexicana</i> (Mez) Kosterm.	90	6.75
<i>Clethra macrophylla</i> DC.	90	6.75
<i>Carpinus tropicalis</i> Walter	214	16.05
Number of individuals with the highest frequency	934	70.07
Total number of individuals (N)	1333	-
CAE		
<i>Inga punctata</i> Willd.	42	1.43
<i>Quercus sapotifolia</i> Liebm.	42	1.43
<i>Alchornea latifolia</i> Sw.	46	1.56
<i>Acrocarpus fraxinifolius</i> Wright & Arn.	47	1.59
<i>Erythrina poeppigiana</i> (Walp.) Skeels.	47	1.59
<i>Eriobotrya japonica</i> Lindley	78	2.65
<i>Enterolobium cyclocarpum</i> (Jacq.) Griseb.	80	2.71
<i>Heliocarpus donnell-smithii</i> Rose	82	2.78
<i>Mimosa scabrella</i> Benth.	144	4.89
<i>Trema micrantha</i> (L.) Blume	145	4.92
<i>I. jinicuil</i> Schltr.	162	5.50
<i>I. latibracteata</i> Harms	163	5.53
<i>Citrus</i> spp.	225	7.63
<i>I. vera</i> Willd.	983	33.36
Number of individuals with the highest frequency	2286	77.57
Total number of individuals (N)	2947	-

Table 2. Diversity components, and indices of proportional abundance of dominance (λ and DS) and equity (Shannon-Wiener) of the three study sites from the coffee region of Coatepec-Xico, Veracruz, Mexico: *i*) undisturbed montane cloud forest (MCF1); *ii*) perturbed montane cloud forest (MCF2), and *iii*) coffee agro-ecosystem (CAE)

	MCF1	MCF2	CAE
Total number of individuals (N)	1388	1333	2947
Species number (S)	252	64	110
Margalef index DMg	34.69	8.756	13.64
Menhinick index DMn	6.764	1.753	2.026
Simpson index λ	0.0075	0.0532	0.1326
Diversity based on Gini–Simpson index ($1-\lambda$)	0.9925	0.9468	0.8674
Shannon-Wiener index	5.1521	3.4613	3.0364

the impacts of land-use change and preserve local biodiversity.

We found that vegetation in CAE included a wide variability of species, and richness increased probably for a species recombination with the MCF surrounding CAE (Villavicencio-Enriquez & Valdez-Hernández, 2003). We observed evidence of this recombination finding species similarities between CAE and the perturbed MCF2. High diversity might be due to a species shift with the MCF nearby (Williams-Linera, 2002). Also, the highest floristic similarity between MCF2 and CAE indicates that CAE is also a perturbed ecosystem. In CAE, the lower diversity in comparison with MCF1 is probably caused by the dominance of some species, partially *Citrus* spp. and *I. vera*.

The floristic composition in CAE is the result of the system's function directed to coffee cultivation. We found in CAE that 33.36% of the individuals were *I. vera*, which are promoted by farmers. Here, it is clear that diversity is influenced by local management, and not only by topography, precipitation or temperature. Trees provide numerous benefits such as building materials, food and firewood, generate family income, promote ecological conditions for wildlife habitats and ecological balance, and also protect against soil erosion (Salam *et al.*, 2000). It has been shown that farmers in agroforestry systems select and eliminate certain tree species according to their preferences and beliefs (Salam *et al.*, 2000; Russell & Franzel, 2004), and also to morphological characteristics (Schroth, 1995); therefore, species composition is conformed by ecological and biophysical variables, and management as well. Also, farmers are paid to modify their farming practice to provide environmental benefits (Salam *et al.*, 2000; Kleijn & Sutherland, 2003). This management provides economic profit and income for local farmers, but it also contributes to improve social levels through the production of important goods, including export crops, fruits, raw material and firewood. Agroforestry systems success lies in the ecological productive capacity over

the long term and also in economic benefits (Michon & de Foresta, 1995).

Maintaining biological diversity is essential for productive agriculture, and ecologically sustainable agriculture is in turn essential for maintaining biological diversity (Pimentel *et al.*, 1992). This maintenance by CAE is reflected in the high number of different species compared to MCF2 and MCF1, which shows CAE's conservation potential, in spite of the presence of exotic (e.g. *Citrus* spp.) and secondary tree species (Table 1) that would be indicators of disturbance. Also, the high proportion of species registered in CAE can support evidence of the services that can provide this system, although we did not evaluate environmental services. Preservation of this agro-ecosystem might represent a possible solution to minimize local biodiversity loss and improve conservation in the central region of Veracruz, especially because the coffee cultivation is more beneficial to the environment than pasture and monocrops such as sugarcane (Esperón-Rodríguez *et al.*, 2016), because coffee conserves tree cover and allows connectivity between open landscapes and forest fragments.

CAE's in Veracruz have a potential role as reservoirs of biodiversity maintaining the forest cover; their conservation as refuges must be considered a priority especially in areas where deforestation and land-use change are increasing. Conservation plans should be addressed to maintain local connection and species recombination between CAE and preserved forests. Knowing the local biodiversity can help local farmers to make better management decision, introducing agroforestry systems with consideration of the markets and products, and also the potential productivity gains and food crops. It must be noticed that although similar, CAE's are not substitute for natural forests; therefore, surrounding forest play an important role in conservation, especially for species that cannot thrive in human modified landscapes. Local management must prioritize the biodiversity preservation and conservation.

Table 3. Tree species diversity comparison among the study sites from the coffee region of Coatepec-Xico, Veracruz, Mexico: *i*) undisturbed montane cloud forest (MCF1); *ii*) perturbed montane cloud forest (MCF2), and *iii*) coffee agro-ecosystem (CAE)

Species	Number of individuals		Percentage (%)	
	MCF1	MCF2	MCF1	MCF2
<i>Alchornea latifolia</i>	17	2	0.1848	0.0054
<i>Cinnamomum effusum</i>	9	13	0.0978	0.0352
<i>Clethra macrophylla</i>	12	90	0.1304	0.2439
<i>Cojoba arborea</i>	1	4	0.0109	0.0108
<i>Liquidambar styraciflua</i>	1	47	0.0109	0.1274
<i>Meliosma alba</i>	2	16	0.0217	0.0434
<i>Myrsine coriacea</i>	2	46	0.0217	0.1247
<i>Ocotea psychotrioides</i>	6	2	0.0652	0.0054
<i>Oreopanax xalapensis</i>	12	4	0.1304	0.0108
<i>Psychotria</i> spp.	17	3	0.1848	0.0081
<i>Quercus xalapensis</i>	3	38	0.0326	0.1029
<i>Styrax glabrescens</i>	4	47	0.0435	0.1274
<i>Symplocos coccinea</i>	1	4	0.0109	0.0108
<i>Trophis mexicana</i>	4	2	0.0435	0.0054
Total number of individuals	92	369	-	-
	MCF2	CAE	MCF2	CAE
<i>Alchornea latifolia</i>	2	46	0.0047	0.0765
<i>Carpinus tropicalis</i>	214	1	0.50352	0.0017
<i>Cinnamomum effusum</i>	13	2	0.0306	0.0033
<i>Citrus</i> spp.	51	225	0.12	0.3744
<i>Cojoba arborea</i>	4	5	0.0094	0.0083
<i>Erythrina americana</i>	5	13	0.0118	0.0216
<i>Heliocarpus donnell-smithii</i>	5	82	0.0118	0.1364
<i>Juglans pyriformis</i>	3	2	0.0071	0.0033
<i>Leucaena leucocephala</i>	2	4	0.0047	0.0067
<i>Lippia myriocephala</i>	11	1	0.0259	0.0017
<i>Myrsine coriacea</i>	46	8	0.1082	0.0133
<i>Quercus sartorii</i>	21	37	0.0494	0.0616
<i>Solanum schlechtendalianum</i>	4	2	0.0094	0.0033
<i>Tapirira mexicana</i>	18	14	0.0424	0.0233
<i>Trema micrantha</i>	3	145	0.0071	0.2413
Unidentified	23	14	0.0541	0.0233
Total number of individuals	425	601	-	-
	MCF1	CAE	MCF	CAE
<i>Alchornea latifolia</i>	17	46	0.3269	0.1411
<i>Cinnamomum effusum</i>	9	2	0.1731	0.0061
<i>Cojoba arborea</i>	1	5	0.0192	0.0153
<i>Dendropanax arboreus</i>	2	9	0.0385	0.0276
<i>Magnolia schiedeana</i>	3	1	0.0577	0.0031
<i>Myrsine coriacea</i>	2	8	0.0385	0.0245
<i>Oreopanax capitatus</i>	1	2	0.0192	0.0061
<i>Oreopanax liebmanni</i>	6	13	0.1154	0.0399
<i>Persea americana</i>	4	7	0.0769	0.0215
<i>Picramnia antidesma</i>	1	7	0.0192	0.0215
<i>Piper nudum</i>	5	1	0.0962	0.0031
Total number of individuals	52	326	-	-

But, what causes this biodiversity in Veracruz? We registered high biodiversity in our study sites despite the size of the sampled area (1500 m²). When we compared our results with previous studies from Veracruz,

Mexico and South America (Table 5) we found a high biodiversity in Veracruz. Although differences may be due to several factors, precipitation is a factor that caught our attention because rainfall is a highly varying

Table 4. Floristic similarity (dissimilarity) components. Indices of similarity/dissimilarity with qualitative and quantitative data of the three study sites from the coffee region of Coatepec-Xico, Veracruz, Mexico: *i*) undisturbed montane cloud forest (MCF1); *ii*) perturbed montane cloud forest (MCF2), and *iii*) coffee agro-ecosystem (CAE)

	MCF1 & MCF2	MCF1 & CAE	MCF2 & CAE
Number of species shared between sites	15	12	16
Indices of similarity/dissimilarity with qualitative data			
Jaccard similarity coefficient IJ	0.0498	0.0343	0.1013
IJ %	4.9834	3.429	10.1266
Sørensen similarity coefficient of IS	0.0949	0.0663	0.1839
IS %	9.4937	6.629	18.3908
Indices of similarity/dissimilarity with quantitative data			
Morisita-Horn index IM-H	0.1796	0.0912	0.1897
IM-H %	17.959	9.115	18.971
Sørensen index (coefficiente of similarity-quantitative)	0.0360	0.0179	0.0640
Iscuant			
Iscuant %	3.602	1.799	6.402

Table 5. Biodiversity studies in montane cloud forests and coffee agro-ecosystems comparing mean annual precipitation (Pp), and species and individual numbers. In bold is indicated data from this work

Location	Pp (mm)	Species	Individuals	Reference
Montane cloud forest				
Teocelo, Veracruz	1500 - 2500	277	600	Luna <i>et al.</i> (1988)
Cofre de Perote, Cortadura, Veracruz	2500	258	Not reported	García <i>et al.</i> (2008)
El Cielo, Tamaulipas	2000	51	2322	Rivas <i>et al.</i> (2005)
Coatepec and Huatusco, Veracruz	1900 - 2000	62	775	López-Gómez <i>et al.</i> (2008)
Central region of Veracruz	1500-2000	83	1029	Williams-Linera (2007)
Central Cordillera of the Colombian Andes	2435	56	Not reported	Cavelier & Tobler (1998)
Western Andean, Peru	1750-2000	88	Not reported	Ledo <i>et al.</i> (2012)
Andean Slope of Bolivia	3500	73	Not reported	Kessler (1999)
Coatepec-Xico, Veracruz	1000-1500	154	2721	-
Coffee agro-ecosystem				
Central region of Veracruz	1500-2000	107	2863	López-Gómez <i>et al.</i> (2008)
Coatepec and Huatusco, Veracruz	1900 - 2000	150	Not reported	Travieso-Bello & Ros (2011)
Coatepec and Huatusco, Veracruz	1900 - 2000	107	2833	Williams-Linera & López-Gómez (2008)
Jitotol, Chiapas	1200 - 3000	50	Not reported	Peeters <i>et al.</i> (2003)
Coatepec-Xico, Veracruz	1000-1500	64	2947	-

parameter in Veracruz (Barradas *et al.*, 2010), and previous studies have shown the importance of water for the species development in the MCF of Veracruz (Esperón-Rodríguez & Barradas, 2015). We hypothesize that the high biodiversity found in our study might

be because of a relatively low precipitation compared to other regions.

Regardless what is causing this high biodiversity, our results indicate that the central region of Veracruz is an important refuge for species, where CAE's

plays an important role in the conservation of biodiversity.

Acknowledgments

We thank to the anonymous reviewers for their critical observations and thoughtful contributions to improve this work.

References

- Albrecht A, Kandji ST, 2003. Carbon sequestration in tropical agroforestry systems. *Agr Ecosyst Environ* 99: 15-27. [http://dx.doi.org/10.1016/S0167-8809\(03\)00138-5](http://dx.doi.org/10.1016/S0167-8809(03)00138-5).
- Aldrich M, Billington NC, Edwards M, Laidlaw R, 1997. Tropical Montane Cloud Forests: An Urgent Priority for Conservation. World Conservation Monitoring Centre, Cambridge, United Kingdom. 407 pp.
- Ashley R, Russell D, Swallow B, 2006. The policy terrain in protected area landscapes: challenges for agroforestry in integrated landscape conservation. *Biodivers Conserv* 15: 663-689. <http://dx.doi.org/10.1007/s10531-005-2100-x>.
- Barradas VL, Cervantes-Pérez J, Ramos-Palacios R, Puchet-Anyul C, Vázquez-Rodríguez P, Granados-Ramírez R, 2010. Meso-scale climate change in the central mountain region of Veracruz State, Mexico. In: Tropical montane cloud forests: science for conservation and management (Brujinzeel LA, Scatena FN, Hamilton LS, eds). Cambridge University Press, Cambridge, UK. pp: 549-556.
- Bellemare J, Motzkin G, Foster DR, 2002. Legacies of the agricultural past in the forested present: an assessment of historical land-use effects on rich mesic forests. *J Biogeogr* 29: 1401-1420. <http://dx.doi.org/10.1046/j.1365-2699.2002.00762.x>.
- Bhagwat SA, Willis KJ, Birks HJB, Whittaker RJ, 2008. Agroforestry: a refuge for tropical biodiversity? *Trends Ecol Evol* 23: 261-264. <http://dx.doi.org/10.1016/j.tree.2008.01.005>.
- Calderón-Aguilera LE, Rivera Monroy V, Porter Bolland L, Martínez Yrizar A, Ladah LB, Martínez Ramos M, Alcocer Alcocer J, Santiago Pérez AL, Hernández Arana H, Reyes Gómez VM, et al., 2012. An assessment of natural and human disturbance effects on Mexican ecosystems: current trends and research gaps. *Biodivers Conserv* 21: 589-617.
- Castillo-Campos G, Reyes S, Abreo ME, 2011. La biodiversidad en Veracruz: Estudio de Estado. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, SECCIÓN IV, Diversidad de Ambientes. Flora y vegetación. Universidad Veracruzana, Instituto de Ecología, A.C. México. 1220 pp.
- Cavelier J, Tobler A, 1998. The effect of abandoned plantations of *Pinus patula* and *Cupressus lusitanica* on soils and regeneration of a tropical montane rain forest in Colombia. *Biodivers Conserv* 7: 335-347. <http://dx.doi.org/10.1023/A:1008829728564>.
- Chapin FS III, Zavaleta ES, Eviner VT, Naylor RL, Vitousek PM, Reynolds HL, Hooper DU, Lavorel S, Sala OE, Honnie SE, Mack MC, Diaz S, 2000. Consequences of changing biodiversity. *Nature* 405: 234-242. <http://dx.doi.org/10.1038/35012241>.
- CONAFOR, 2011. Silvicultura comunitaria. Available at: www.conafor.gob.mx/portal/index.php/temas-forestales/silvicultura, Accessed November, 2014.
- Cronquist A, 1988. The Evolution and Classification of Flowering Plants. The New York Botanical Garden, N.Y. 555 pp.
- Current D, Lutz E, Scherr SJ, (eds), 1995. Costs, benefits, and farmer adoption of agroforestry: project experience in Central America and the Caribbean. World Bank Environment Paper No. 14. Washington D.C. 212 pp. <http://dx.doi.org/10.1596/0-8213-3428-X>.
- Dávalos Sotelo R, Morato MI, Martínez Pinillos-Cueto E, 2008. Almacenamiento de carbono. In: Agroecosistemas cafetaleros de Veracruz: biodiversidad, manejo y conservación (Manson RH, Hernández-Ortiz V, Gallina S, Mehlreter K, eds.). Instituto de Ecología A.C. (INECOL), Instituto Nacional de Ecología (INE-SEMARNAT), México, pp: 223-234.
- Ellis E, Martínez M, 2010. Vegetación y Uso de Suelo de Veracruz. In: Atlas del Patrimonio Natural, Histórico y Cultural del Estado de Veracruz (Tomo 1, Patrimonio Natural). Comisión para la Conmemoración del Bicentenario de la Independencia Nacional y del Centenario de la Revolución Mexicana. Gobierno del Estado de Veracruz. Xalapa, Veracruz, pp: 203-226.
- Esperón-Rodríguez M, Barradas VL, 2015. Comparing environmental vulnerability in the montane cloud forest of eastern Mexico: A vulnerability index. *Ecol Indic* 52: 300-310. <http://dx.doi.org/10.1016/j.ecolind.2014.12.019>.
- Esperón-Rodríguez M, Bonifacio-Martínez M, Barradas VL, 2016. Socio-economical vulnerability to climate change in the central mountainous region of eastern Mexico. *Ambio* 45:146-160. <http://dx.doi.org/10.1007/s13280-015-0690-4>.
- García JG, Castillo G, Mehlreter K, Martínez ML, Vázquez G, 2008. Composición florística de un bosque mesófilo de montaña del centro de Veracruz, México. *Bol Soc Bot Méx* 83: 37-52.
- Geissert D, Ibáñez A, 2008. Calidad y ambiente físico-químico de los suelos. In: Agroecosistemas cafetaleros de Veracruz: biodiversidad, manejo y conservación (Manson RH, Hernández-Ortiz V, Gallina S, Mehlreter K, eds.). Instituto de Ecología A.C. (INECOL), Instituto Nacional de Ecología (INE-SEMARNAT), México, pp: 213-222.
- Gómez-Pompa A, 1978. Ecología de la vegetación del estado de Veracruz. CECSA. México. 91 pp.
- González-Espinosa M, Meave JA, Ramírez-Marcial N, Toledo-Aceves T, Lorea-Hernández, FG, Ibarra-Manríquez G, 2012. Los bosques de niebla de México: conservación y restauración de su componente arbóreo. *Ecosistemas* 21: 36-54.
- Kessler M, 1999. Plant species richness and endemism during natural landslide succession in a perhumid montane forest in the Bolivian Andes. *Ecotropica* 5: 123-136.

- Kleijn D, Sutherland JW, 2003. How effective are European agro-environment schemes in conserving and promoting biodiversity? *J Appl Ecol* 40: 947-969. <http://dx.doi.org/10.1111/j.1365-2664.2003.00868.x>.
- Landeros-Sánchez C, Moreno-Seceña JC, Escamilla-Prado E, Ruiz-Bello R, 2011. La biodiversidad en Veracruz: Estudio de Estado. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, SECCIÓN VII, Sistemas productivos y alternativas económicas sustentables. Diversificación de cultivos. Universidad Veracruzana, Instituto de Ecología, A.C. México, pp: 463-475.
- Ledo A, Condés S, Alberdi I, 2012. Forest biodiversity assessment in Peruvian Andean montane cloud forest. *J Mount Sci* 9: 372-384. <http://dx.doi.org/10.1007/s11629-009-2172-2>.
- López-Gómez AM, Williams-Linera G, Manson RH, 2008. Tree species diversity and vegetation structure in shade coffee farms in Veracruz, Mexico. *Agr Ecosyst Environ* 124: 160-172. <http://dx.doi.org/10.1016/j.agee.2007.09.008>.
- Luna I, Almeida L, Villers L, Lorenzo L, 1988. Reconocimiento florístico y consideraciones fitogeográficas del bosque mesófilo de montaña de Teocelo, Veracruz. *Bol Soc Bot Méx* 48: 35-63.
- Manson RH, Hernández-Ortiz V, Gallina S, Mehltreter K (eds.), 2008. Agroecosistemas cafetaleros de Veracruz: biodiversidad, manejo y conservación. Instituto de Ecología A.C. (INECOL), Instituto Nacional de Ecología (INE-SEMARNAT), México. 330 pp.
- Mcneely JA, 1995. How traditional agro-ecosystems can contribute to conserving biodiversity. In: *Conserving biodiversity outside protected areas. The role of traditional agro-ecosystems* (Halladay P, Gilmour DA, eds.). IUCN, Gland, Switzerland, and Cambridge, UK in collaboration with AMA Andalucía and Centro de Investigación F. González-Bernáldez, pp: 20-40.
- MEA, 2005. *Ecosystems and Human well-being. Millenium Ecosystem Assessment*, Island Press, New York.
- Mejía NR, Meave JA, Ruiz CA, 2004. Análisis estructural de un bosque mesófilo de montaña en el extremo oriental de la sierra madre del sur (Oaxaca), México. *Bol Soc Bot Méx* 74: 13-29.
- Michon G, De Foresta H, 1995 The Indonesian agro-forest model. *Conserving biodiversity outside protected areas. In: The role of traditional agro-ecosystems* (Halladay P, Gilmour DA, eds.). IUCN, Gland, Switzerland, and Cambridge, UK in collaboration with AMA Andalucía and Centro de Investigación F. González-Bernáldez, pp: 90-104.
- Moreno CE, 2001. Métodos para medir la biodiversidad. *M&T-Manuales y Tesis SEA*, vol. 1. Zaragoza, Spain.
- Mostacedo B, Fredericksen TS, 2000. *Manual de Métodos Básicos de Muestreo y Análisis en Ecología Vegetal*. Santa Cruz, Bolivia. BOLFOR. 87 pp.
- Nonato De Souza H, De Goede RGM, Brussaard L, Cardoso IM, Duarte EMG, F RBA, Gomes LC, Pulleman MM, 2012. Protective shade, tree diversity and soil properties in coffee agroforestry systems in the Atlantic Rainforest biome. *Agr Ecosys Environ* 146: 179-196. <http://dx.doi.org/10.1016/j.agee.2011.11.007>.
- O'Dea N, Whittaker RJ, 2007. How resilient are Andean montane forest bird communities to habitat degradation? *Biodivers Conserv* 16: 1131-1159. <http://dx.doi.org/10.1007/s10531-006-9095-9>.
- Olguín J, Sánchez- Galván G, Vidal G, 2011. La biodiversidad en Veracruz: Estudio de Estado. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, SECCIÓN V, La biodiversidad del estado y algunas de sus amenazas . La producción de café como amenaza a la biodiversidad. Universidad Veracruzana, Instituto de Ecología, A.C. México, pp: 391-425.
- Ortega F, Castillo, G, 1996. El bosque mesófilo de montaña y sus implicaciones forestales. *Ciencias* 43: 32-39.
- Palacio-Prieto JL, Bocco G, Velásquez A, Mas JF, Takaki-Takaki F, Victoria A, Luna González L, Gómez-Rodríguez G, López-García J, Palma M, et al., 2000. Technical Note: Current situation of forest resources in Mexico: results of the 2000 National Forest Inventory. *Inv Geogr Bol Inst Geogr* 43: 183-203.
- Peeters L, Soto-Pinto L, Perales H, Montoya G, Ishiki M, 2003. Coffee production, timber, and redwood in traditional and Inga-shaded plantations in Southern Mexico. *Agr Ecosys Environ* 95: 481-493. [http://dx.doi.org/10.1016/S0167-8809\(02\)00204-9](http://dx.doi.org/10.1016/S0167-8809(02)00204-9).
- Pimentel D, Stachow U, Takacs DA, Brubaker HW, Dumas AR, Meaney JJ, O'Neil J, Onsi DE, Corzilius DB, 1992. *Conserving Biological Diversity in Agricultural Forestry Systems -Most Biological Diversity Exists in Human-Managed Ecosystems*. *Bioscience* 42(5): 354-362. <http://dx.doi.org/10.2307/1311782>.
- R Core Team, 2014. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rivas CJ, Aguirre C, Jiménez P, Corral R, 2005. Un análisis del efecto del aprovechamiento forestal sobre la diversidad estructural en el bosque mesófilo de montaña «El Cielo», Tamaulipas, México. *Sist Rec Forest* 14(2): 217-228.
- Ruelas Monjardín LC, Nava Tablada ME, Cervantes J, Barradas VL, 2014. Importancia ambiental de los agroecosistemas cafetaleros bajo sombra en la zona central montañosa del estado de Veracruz, México. *Madera y Bosque* 20: 27-40.
- Russell D, Franzel S, 2004. Trees of prosperity: Agroforestry, markets and the African smallholder. *Agroforest Syst* 61: 345-355. <http://dx.doi.org/10.1023/B:AGFO.0000029009.53337.33>.
- SAGARPA, 2007. *Sistema Integral in Información Agroalimentaria y Pesquera (SIAP)*. Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimento de México. (<http://www.siap.sagarpa.gob.mx/>).
- Sala OE, Chapin FS III, Armesto JJ, Berlow R, Bloomfield J, Dirzo R, Mooney HA, Oesterheld M, Poff NL, Sykes MT, Walker BH, Walker W, Wall DH, 2000. Biodiversity – global biodiversity scenarios for the year 2100. *Science* 287: 1770-1774. <http://dx.doi.org/10.1126/science.287.5459.1770>.

- Salam MA, Noguchi T, Koike M, 2000. Understanding why farmers plant trees in the homestead agroforestry in Bangladesh. *Agroforest Syst* 50: 77-93. <http://dx.doi.org/10.1023/A:1006403101782>.
- Schroth G, 1995. Tree root characteristics as criteria for species selection and systems design in agroforestry. *Agroforest Syst* 30: 125-143. <http://dx.doi.org/10.1007/BF00708917>.
- Sosa V, Gómez-Pompa A (eds.), 1994. Lista Florística. In *Flora de Veracruz*. Instituto de Ecología, A.C., Xalapa, Veracruz, México. 245 pp.
- Thangata PH, Hildebrand PE, 2012. Carbon stock and sequestration potential of agroforestry systems in smallholder agroecosystems of sub-Saharan Africa: Mechanisms for reducing emissions from deforestation and forest degradation (REDD+). *Agr Ecosys Environ* 158: 172-183. <http://dx.doi.org/10.1016/j.agee.2012.06.007>.
- Tilman D, Cassman KG, Matson PA, Naylor R, Polasky S, 2002. Agricultural sustainability and intensive production practices. *Nature* 418: 671-677. <http://dx.doi.org/10.1038/nature01014>.
- Toledo-Aceves T, Meave JA, González-Espinosa M, Ramírez-Marcial N, 2011. Tropical montane cloud forests: current threats and opportunities for their conservation and sustainable management in Mexico. *J Environ Manag* 92(3): 974-981. <http://dx.doi.org/10.1016/j.jenvman.2010.11.007>.
- Tornquist CG, Hons FM, Feagley SE, Haggard J, 1999. Agroforestry system effects on soil characteristics of the Sarapiquí region of Costa Rica. *Agr Ecos Environ* 73: 19-28. [http://dx.doi.org/10.1016/S0167-8809\(99\)00004-3](http://dx.doi.org/10.1016/S0167-8809(99)00004-3).
- Travieso-Bello AC, Ros AV, 2011. La biodiversidad en Veracruz: Estudio de Estado. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, Gobierno del Estado de Veracruz, Universidad Veracruzana, Instituto de Ecología, A.C. México, pp: 261-270.
- Villarreal H, Álvarez M, Córdoba S, Escobar F, Fagua G, Gast F, Mendoza H, Ospina M, Umaña AM, 2004. Manual de métodos para el desarrollo de inventarios de biodiversidad. Programa de inventarios de Biodiversidad. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Bogotá, Colombia. pp: 185-227.
- Villavicencio-Enríquez L, Valdez-Hernández JI, 2003. Análisis de la estructura arbórea del sistema agroforestal rusticano de café en San Miguel, Veracruz, México. *Agrociencia* 37(4): 413-423.
- Williams-Linera G, 2002. Tree species richness complementarity, disturbance and fragmentation in a Mexican tropical montane cloud forest. *Biodivers Conserv* 11: 1825-1843. <http://dx.doi.org/10.1023/A:1020346519085>.
- Williams-Linera G, Manson RH, Isonza-Vera E, 2002. La fragmentación del bosque mesófilo de montaña y patrones de uso del suelo en la región oeste de Xalapa, Veracruz, México. *Madera y Bosques* 8:73-89.
- Williams-Linera G, 2007. El bosque de niebla del centro de Veracruz: ecología, historia y destino en tiempos de fragmentación y cambio climático. CONABIO, Instituto de Ecología, A.C. México. 197 pp.
- Williams-Linera G, López-Gómez A, 2008. Estructura y diversidad de la vegetación leñosa. In: *Agroecosistemas cafetaleros de Veracruz: biodiversidad, manejo y conservación* (Manson RH, Hernández-Ortiz V, Gallina S, Melhlreter K, eds.). Instituto de Ecología A.C. and Instituto Nacional de Ecología (INE-SEMARNAT), México, pp. 55-63.
- Wright SJ, 2005. Tropical forests in a changing environment. *Trends Ecol Evol* 20:5 53-60.

[Log in to My Ulrich's](#)Macquarie University Library —Select Language—[Search](#) [Workspace](#) [Ulrich's Update](#) [Admin](#)

Enter a Title, ISSN, or search term to find journals or other periodicals:

2171-9845

[▶ Advanced Search](#)[Search Results](#)JCR® Web [Search MultiSearch: \[ISSN Search\] \[Title Search\]](#)

Forest Systems

Title Details

Related Titles

[▶ Alternative Media Edition \(1\)](#)

Lists

[Marked Titles \(0\)](#)

Search History

[2171-5068 - \(1\)](#)[2171-9845 - \(1\)](#)[Save to List](#) [Email](#) [Download](#) [Print](#) [Corrections](#) [Expand All](#) [Collapse All](#)

▼ Basic Description

Title	Forest Systems
ISSN	2171-9845
Publisher	Instituto Nacional de Investigacion y Tecnologia Agraria y Alimentaria (I N I A)
Country	Spain
Status	Active
Frequency	3 times a year
Language of Text	Text in: English
Refereed	Yes
Abstracted / Indexed	Yes
Open Access	Yes http://recyt.fecyt.es/index.php/IA/index
Serial Type	Journal
Content Type	Academic / Scholarly
Format	Online
Website	http://recyt.fecyt.es/index.php/IA/index

▶ Subject Classifications

▶ Additional Title Details

▶ Title History Details

▶ Publisher & Ordering Details

▶ Online Availability

▶ Abstracting & Indexing

▶ Demographics

[Save to List](#) [Email](#) [Download](#) [Print](#) [Corrections](#) [Expand All](#) [Collapse All](#)