

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368


Editors: W. Clegg and D. G. Watson

## 2,8-Dibromo-4,10-dichloro-6*H*,12*H*-5,11-methano-dibenzo[*b,f*][1,5]diazocine

Kai-Xian Zhu, Donald C. Craig and Andrew C. Try

*Acta Cryst.* (2008). E64, o1797

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



*Acta Crystallographica Section E: Structure Reports Online* is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals **Online** is available from [journals.iucr.org](http://journals.iucr.org)

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

2,8-Dibromo-4,10-dichloro-6*H*,12*H*-5,11-methanodibenzo[*b,f*][1,5]diazocineKai-Xian Zhu,<sup>a</sup> Donald C. Craig<sup>b</sup> and Andrew C. Try<sup>a\*</sup><sup>a</sup>Department of Chemistry and Biomolecular Sciences, Building F7B, Macquarie University, Sydney, NSW 2109, Australia, and <sup>b</sup>School of Chemistry, University of New South Wales, Sydney, NSW 2052, Australia

Correspondence e-mail: andrew.try@mq.edu.au


Received 31 July 2008; accepted 14 August 2008

Key indicators: single-crystal X-ray study;  $T = 294$  K; mean  $\sigma(\text{C}-\text{C}) = 0.020$  Å;  $R$  factor = 0.057;  $wR$  factor = 0.062; data-to-parameter ratio = 7.4.

The title compound,  $\text{C}_{15}\text{H}_{10}\text{Br}_2\text{Cl}_2\text{N}_2$ , a 2,8-dibromo-4,10-dichloro Tröger's base analogue derived from 4-bromo-2-chloroaniline, has a dihedral angle of  $110.9(10)^\circ$  between the two aryl rings, the largest yet measured for a simple dibenzo analogue.

## Related literature

For related literature on the synthesis and crystal structures of dihalogenated Tröger's base analogues, see: Jensen & Wärnmark (2001); Faroughi *et al.* (2006*a*, 2007*a,b*). For Tröger's base analogues substituted with multiple electron-withdrawing groups, see: Faroughi *et al.* (2006*b*); Bhuiyan *et al.* (2006, 2007); Vande Velde *et al.* (2008). For reactions of halogenated Tröger's base analogues, see: Jensen *et al.* (2002); Hof *et al.* (2005). For literature on racemization of Tröger's base analogues and the effect of substituents *ortho* to the diazocine N atoms, see: Lenev *et al.* (2006).



## Experimental

## Crystal data

 $\text{C}_{15}\text{H}_{10}\text{Br}_2\text{Cl}_2\text{N}_2$   
 $M_r = 449.0$ Orthorhombic,  $Pca2_1$   
 $a = 7.910(2)$  Å $b = 12.601(3)$  Å  
 $c = 15.230(4)$  Å  
 $V = 1518.0(7)$  Å<sup>3</sup>  
 $Z = 4$ Mo  $K\alpha$  radiation  
 $\mu = 5.64$  mm<sup>-1</sup>  
 $T = 294$  K  
 $0.30 \times 0.12 \times 0.07$  mm

## Data collection

Enraf–Nonius CAD-4  
diffractometer  
Absorption correction: analytical  
(de Meulenaer & Tompa, 1965)  
 $T_{\min} = 0.52$ ,  $T_{\max} = 0.69$   
1394 measured reflections1394 independent reflections  
1028 reflections with  $I > 2\sigma(I)$   
1 standard reflection  
frequency: 30 min  
intensity decay: none

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.056$   
 $wR(F^2) = 0.061$   
 $S = 1.61$   
1394 reflections  
189 parametersH-atom parameters constrained  
 $\Delta\rho_{\max} = 0.98$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -1.02$  e Å<sup>-3</sup>  
Absolute structure: Flack (1983)  
Flack parameter: 0.09 (2)

Data collection: *CAD-4* (Schagen *et al.*, 1989); cell refinement: *CAD-4*; data reduction: local program; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *RAELS* (Rae, 1996); molecular graphics: *ORTEPII* (Johnson, 1976); software used to prepare material for publication: local programs.

The authors thank Macquarie University for the award of a Macquarie University Research Development Grant to ACT.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2290).

## References

- Altomare, A., Casciarano, G. L., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.  
Bhuiyan, M. D. H., Jensen, P. & Try, A. C. (2007). *Acta Cryst.* **E63**, o4393.  
Bhuiyan, M. D. H., Try, A. C., Klepetko, J. & Turner, P. (2006). *Acta Cryst.* **E62**, o4887–o4888.  
Faroughi, M., Try, A. C., Klepetko, J. & Turner, P. (2007*a*). *Tetrahedron Lett.* **48**, 6548–6551.  
Faroughi, M., Try, A. C. & Turner, P. (2006*a*). *Acta Cryst.* **E62**, o3674–o3675.  
Faroughi, M., Try, A. C. & Turner, P. (2006*b*). *Acta Cryst.* **E62**, o3893–o3894.  
Faroughi, M., Try, A. C. & Turner, P. (2007*b*). *Acta Cryst.* **E63**, o2695.  
Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.  
Hof, F., Schar, M., Scofield, D. M., Fischer, F., Diederich, F. & Sergeev, S. (2005). *Helv. Chim. Acta*, **88**, 2333–2344.  
Jensen, J., Strozyk, M. & Wärnmark, K. (2002). *Synthesis*, pp. 2761–2765.  
Jensen, J. & Wärnmark, K. (2001). *Synthesis*, pp. 1873–1877.  
Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.  
Lenev, D. A., Lyssenko, K. A., Golovanov, D. G., Buss, V. & Kostyanovsky, R. G. (2006). *Chem. Eur. J.* **12**, 6412–6418.  
Meulenaer, J. de & Tompa, H. (1965). *Acta Cryst.* **19**, 1014–1018.  
Rae, A. D. (1996). *RAELS*. University of New South Wales, Australia.  
Schagen, J. D., Straver, L., van Meurs, F. & Williams, G. (1989). *CAD-4 Manual*. Enraf–Nonius, Delft, The Netherlands.  
Vande Velde, C. M. L., Didier, D., Blockhuys, F. & Sergeev, S. (2008). *Acta Cryst.* **E64**, o538.

## **supplementary materials**

*Acta Cryst.* (2008). E64, o1797 [ doi:10.1107/S1600536808026226 ]

## 2,8-Dibromo-4,10-dichloro-6*H*,12*H*-5,11-methanodibenzo[*b,f*][1,5]diazocine

K.-X. Zhu, D. C. Craig and A. C. Try

### Comment

Tröger's base analogues bearing electron-withdrawing groups were long thought to be difficult, if not impossible, to prepare. However, the synthesis of dihalogenated (Jensen & Wärnmark, 2001), octafluoro (Vande Velde *et al.*, 2008) and tetranitro (Bhuiyan *et al.*, 2007) Tröger's base analogues highlight the possibilities that now exist in terms of incorporating electron-withdrawing groups on the starting anilines. The synthetic utility of halogen-substituted Tröger's base analogue has been demonstrated with their conversion to alkyne- (Jensen & Wärnmark, 2001; Jensen *et al.*, 2002) and functionalized phenyl- (Hof *et al.*, 2005) substituted analogues, among others. It is noteworthy that crystal structures of several other 2,4,8,10-tetrasubstituted Tröger's base analogues exhibit large dihedral angles that are close to that in (I). Tröger's base analogues are known to undergo racemization in acidic solution, however the presence of a substituent at the *ortho*-position, relative to the bridge nitrogen atoms, has been shown to increase the racemization barrier (Lenev *et al.*, 2006).

The molecular structure of (I) is shown in Fig. 1 and it was prepared as outlined in Fig. 2.

### Experimental

4-Bromo-2-chloroaniline (1 g, 4.84 mmol) and paraformaldehyde (232 mg, 7.74 mmol) were added to an ice-cold solution of trifluoroacetic acid (10 ml). The reaction mixture was then stirred in dark at room temperature for 7 days under an atmosphere of argon. The ice-cold reaction mixture was basified by the dropwise addition of a mixture of ammonia (28%, 20 ml) and water (40 ml), followed by the addition of a saturated sodium hydrogen carbonate solution (20 ml). The resultant mixture was then extracted with dichloromethane (3 x 20 ml) and the combined organic layers were washed with brine (40 ml), dried over anhydrous sodium sulfate, filtered and evaporated to dryness. The crude product was chromatographed (silica gel, dichloromethane:hexane 8:2) to afford 2,8-dibromo-4,10-dichloro-6*H*,12*H*-5,11-methanodibenzo[*b,f*][1,5]diazocine (I) (613 mg, 56%) as a white solid and as a racemic mixture: m.p. 471–472 K; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 4.21–4.33 (4*H*, m), 4.55 (2*H*, d, *J* 17.3 Hz), 7.04 (2*H*, d, *J* 2.1 Hz), 7.41 (2*H*, d, *J* 2.1 Hz); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 54.37, 67.32, 117.15, 128.49, 130.17, 131.02, 131.71, 142.33. Analysis found: C 40.46; H 2.22; N 6.46; C<sub>15</sub>H<sub>10</sub>Br<sub>2</sub>Cl<sub>2</sub>N<sub>2</sub> requires C 40.13; H 2.25; N 6.24. Single crystals were obtained from slow evaporation from dichloromethane solution of (I).

### Refinement

Hydrogen atoms were included in positions calculated each cycle (C—H = 1.0 Å), and were assigned thermal parameters equal to their bonded atom. The maximum and minimum electron density peaks were located 0.73 and 1.20 Å from the Cl2 and Br1 atoms, respectively.

## Figures




Fig. 1. *ORTEP* (Johnson, 1976) plot of (I), with ellipsoids at the 10% probability level. H atoms are drawn as spheres of arbitrary radius.



Fig. 2. Synthetic scheme for the synthesis of (I) showing the numbering system used in naming the compound.

## 2,8-Dibromo-4,10-dichloro-6*H*,12*H*-5,11-methanodibenzo[*b,f*][1,5]diazocine

### Crystal data

$C_{15}H_{10}Br_2Cl_2N_2$

$M_r = 449.0$

Orthorhombic,  $Pca2_1$

Hall symbol:  $P\ 2c\ -2ac$

$a = 7.910\ (2)\ \text{\AA}$

$b = 12.601\ (3)\ \text{\AA}$

$c = 15.230\ (4)\ \text{\AA}$

$V = 1518.0\ (7)\ \text{\AA}^3$

$Z = 4$

$F_{000} = 872.0$

$D_x = 1.96\ \text{Mg m}^{-3}$

Melting point: 471 K

Mo  $K\alpha$  radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 11 reflections

$\theta = 10\text{--}11^\circ$

$\mu = 5.64\ \text{mm}^{-1}$

$T = 294\ \text{K}$

Prism, colourless

$0.30 \times 0.12 \times 0.07\ \text{mm}$

### Data collection

Enraf–Nonius CAD-4  
diffractometer

$\omega$ - $2\theta$  scans

Absorption correction: analytical  
de Meulenaer & Tompa (1965)

$T_{\min} = 0.52$ ,  $T_{\max} = 0.69$

1394 measured reflections

1394 independent reflections

1028 reflections with  $I > 2\sigma(I)$

$\theta_{\max} = 25^\circ$

$h = 0 \rightarrow 9$

$k = 0 \rightarrow 14$

$l = -18 \rightarrow 0$

1 standard reflections

every 30 min

intensity decay: none

### Refinement

Refinement on  $F$

$R[F^2 > 2\sigma(F^2)] = 0.056$

$wR(F^2) = 0.061$

$S = 1.61$

1394 reflections

189 parameters

H-atom parameters constrained

$w = 1/[\sigma^2(F) + 0.0004F^2]$

$(\Delta/\sigma)_{\max} = 0.002$

$\Delta\rho_{\max} = 0.98\ \text{e \AA}^{-3}$

$\Delta\rho_{\min} = -1.02\ \text{e \AA}^{-3}$

Extinction correction: none

Absolute structure: Flack (1983), 0 Friedel pairs

Flack parameter: 0.09 (2)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.8061 (2)	0.6867 (1)	0.3991 (1)	0.0585 (5)
Br2	0.2324 (2)	0.0800 (1)	0.0428 (2)	0.0527 (5)
Cl1	0.9770 (5)	0.5167 (4)	0.0779 (3)	0.057 (1)
Cl2	0.5654 (5)	0.0565 (3)	0.3603 (3)	0.052 (1)
N1	0.9460 (14)	0.3064 (10)	0.1658 (8)	0.041 (3)
N2	0.8319 (15)	0.1842 (9)	0.2739 (8)	0.044 (3)
C1	0.9798 (19)	0.2123 (12)	0.2198 (10)	0.044 (4)
C2	0.8998 (18)	0.3929 (12)	0.2193 (10)	0.043 (4)
C3	0.8445 (18)	0.3790 (11)	0.3054 (8)	0.041 (4)
C4	0.817 (2)	0.2693 (12)	0.3414 (8)	0.049 (4)
C5	0.6842 (19)	0.1696 (11)	0.2226 (9)	0.039 (4)
C6	0.6740 (19)	0.2086 (11)	0.1360 (9)	0.036 (4)
C7	0.8206 (17)	0.2753 (12)	0.0978 (9)	0.040 (4)
C8	0.9155 (17)	0.4984 (13)	0.1855 (9)	0.043 (4)
C9	0.8825 (19)	0.5861 (12)	0.2397 (11)	0.048 (4)
C10	0.839 (2)	0.5684 (13)	0.3247 (11)	0.050 (4)
C11	0.818 (2)	0.4666 (13)	0.3570 (10)	0.056 (4)
C12	0.5565 (17)	0.1082 (10)	0.2538 (8)	0.031 (3)
C13	0.4158 (17)	0.0854 (10)	0.2036 (8)	0.035 (3)
C14	0.4117 (17)	0.1221 (11)	0.1168 (9)	0.040 (4)
C15	0.5433 (19)	0.1819 (10)	0.0853 (9)	0.034 (3)
H1C1	1.0780	0.2273	0.2593	0.044
H2C1	1.0078	0.1513	0.1804	0.044
H1C4	0.7008	0.2660	0.3674	0.049
H2C4	0.9027	0.2559	0.3883	0.049
H1C7	0.7726	0.3410	0.0708	0.040
H2C7	0.8794	0.2326	0.0517	0.040
HC9	0.8906	0.6600	0.2162	0.048
HC11	0.7825	0.4567	0.4195	0.056
HC13	0.3194	0.0438	0.2286	0.035
HC15	0.5411	0.2061	0.0228	0.034

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.060 (1)	0.060 (1)	0.0549 (9)	-0.0011 (8)	-0.001 (1)	-0.0138 (9)
Br2	0.0470 (9)	0.0650 (9)	0.0461 (8)	-0.0118 (8)	-0.0026 (9)	-0.0091 (9)
Cl1	0.061 (3)	0.066 (3)	0.044 (2)	-0.019 (2)	0.010 (2)	0.000 (2)
Cl2	0.063 (3)	0.056 (2)	0.037 (2)	0.000 (2)	0.006 (2)	0.018 (2)
N1	0.029 (7)	0.058 (8)	0.037 (7)	0.001 (6)	0.002 (6)	-0.003 (6)
N2	0.046 (7)	0.048 (7)	0.038 (7)	0.004 (6)	-0.009 (6)	0.020 (6)
C1	0.029 (9)	0.067 (9)	0.036 (8)	0.005 (8)	-0.016 (7)	0.012 (8)
C2	0.051 (9)	0.043 (9)	0.034 (8)	0.009 (8)	0.020 (8)	-0.006 (7)
C3	0.045 (9)	0.059 (9)	0.019 (8)	-0.012 (8)	-0.004 (7)	-0.005 (7)

## supplementary materials

---

C4	0.083 (9)	0.053 (8)	0.013 (7)	0.004 (9)	-0.006 (7)	0.007 (7)
C5	0.048 (9)	0.043 (9)	0.026 (7)	0.000 (7)	0.013 (7)	0.007 (7)
C6	0.039 (8)	0.042 (8)	0.026 (8)	0.010 (8)	0.002 (6)	-0.006 (7)
C7	0.037 (8)	0.052 (9)	0.030 (8)	-0.007 (8)	0.007 (7)	-0.002 (7)
C8	0.029 (8)	0.062 (9)	0.039 (9)	-0.002 (8)	-0.002 (7)	0.004 (9)
C9	0.047 (9)	0.038 (9)	0.059 (9)	-0.010 (8)	-0.001 (8)	-0.007 (8)
C10	0.047 (9)	0.057 (9)	0.046 (9)	-0.005 (8)	0.017 (8)	-0.007 (8)
C11	0.085 (9)	0.052 (9)	0.029 (8)	-0.011 (9)	0.009 (9)	-0.007 (8)
C12	0.037 (8)	0.032 (8)	0.024 (7)	0.009 (7)	0.010 (6)	-0.002 (6)
C13	0.046 (9)	0.043 (9)	0.016 (6)	0.014 (8)	0.001 (6)	0.000 (7)
C14	0.029 (8)	0.039 (8)	0.052 (9)	0.006 (7)	0.010 (7)	-0.015 (8)
C15	0.034 (8)	0.039 (8)	0.027 (7)	0.004 (7)	0.009 (7)	-0.004 (7)

### *Geometric parameters (Å, °)*

Br1—C10	1.890 (15)	C4—H2C4	1.000
Br2—C14	1.888 (14)	C5—C6	1.410 (18)
C11—C8	1.724 (14)	C5—C12	1.358 (18)
C12—C12	1.750 (13)	C6—C7	1.546 (20)
N1—C1	1.468 (18)	C6—C15	1.333 (19)
N1—C2	1.409 (17)	C7—H1C7	1.000
N1—C7	1.487 (18)	C7—H2C7	1.000
N2—C1	1.474 (19)	C8—C9	1.404 (20)
N2—C4	1.491 (18)	C9—C10	1.357 (19)
N2—C5	1.417 (18)	C9—HC9	1.000
C1—H1C1	1.000	C10—C11	1.385 (21)
C1—H2C1	1.000	C11—HC11	1.000
C2—C3	1.394 (18)	C12—C13	1.381 (17)
C2—C8	1.431 (20)	C13—C14	1.402 (18)
C3—C4	1.503 (20)	C13—HC13	1.000
C3—C11	1.371 (20)	C14—C15	1.371 (19)
C4—H1C4	1.000	C15—HC15	1.000
C1—N1—C2	110.4 (12)	N1—C7—C6	112.5 (11)
C1—N1—C7	107.4 (11)	N1—C7—H1C7	108.7
C2—N1—C7	115.7 (11)	N1—C7—H2C7	108.7
C1—N2—C4	106.0 (11)	C6—C7—H1C7	108.7
C1—N2—C5	112.2 (11)	C6—C7—H2C7	108.7
C4—N2—C5	114.1 (12)	H1C7—C7—H2C7	109.5
N1—C1—N2	111.3 (11)	C11—C8—C2	119.3 (11)
N1—C1—H1C1	109.0	C11—C8—C9	120.4 (12)
N1—C1—H2C1	109.0	C2—C8—C9	120.2 (12)
N2—C1—H1C1	109.0	C8—C9—C10	118.6 (15)
N2—C1—H2C1	109.0	C8—C9—HC9	120.7
H1C1—C1—H2C1	109.5	C10—C9—HC9	120.7
N1—C2—C3	121.9 (14)	Br1—C10—C9	118.5 (13)
N1—C2—C8	119.2 (12)	Br1—C10—C11	120.0 (11)
C3—C2—C8	118.8 (13)	C9—C10—C11	121.4 (15)
C2—C3—C4	120.3 (13)	C3—C11—C10	121.6 (14)
C2—C3—C11	119.1 (14)	C3—C11—HC11	119.2

C4—C3—C11	120.6 (12)	C10—C11—HC11	119.2
N2—C4—C3	113.5 (10)	Cl2—C12—C5	120.5 (12)
N2—C4—H1C4	108.4	Cl2—C12—C13	117.9 (10)
N2—C4—H2C4	108.4	C5—C12—C13	121.6 (13)
C3—C4—H1C4	108.4	C12—C13—C14	118.2 (13)
C3—C4—H2C4	108.4	C12—C13—HC13	120.9
H1C4—C4—H2C4	109.5	C14—C13—HC13	120.9
N2—C5—C6	121.2 (13)	Br2—C14—C13	119.2 (11)
N2—C5—C12	119.6 (13)	Br2—C14—C15	121.0 (11)
C6—C5—C12	118.9 (15)	C13—C14—C15	119.6 (13)
C5—C6—C7	119.8 (13)	C6—C15—C14	121.6 (13)
C5—C6—C15	119.9 (14)	C6—C15—HC15	119.2
C7—C6—C15	120.1 (12)	C14—C15—HC15	119.2
C2—N1—C1—N2	57.5 (15)	C2—C3—C11—C10	2.3 (24)
C2—N1—C1—H1C1	-62.8	C2—C3—C11—HC11	-177.7
C2—N1—C1—H2C1	177.8	C4—C3—C11—C10	-178.6 (16)
C7—N1—C1—N2	-69.4 (15)	C4—C3—C11—HC11	1.4
C7—N1—C1—H1C1	170.3	N2—C5—C6—C7	-4.4 (20)
C7—N1—C1—H2C1	50.8	N2—C5—C6—C15	171.2 (13)
C1—N1—C2—C3	-18.2 (18)	C12—C5—C6—C7	-177.9 (12)
C1—N1—C2—C8	159.5 (13)	C12—C5—C6—C15	-2.3 (20)
C7—N1—C2—C3	103.9 (16)	N2—C5—C12—Cl2	5.4 (18)
C7—N1—C2—C8	-78.3 (17)	N2—C5—C12—C13	-175.2 (12)
C1—N1—C7—C6	44.6 (15)	C6—C5—C12—Cl2	179.1 (10)
C1—N1—C7—H1C7	165.1	C6—C5—C12—C13	-1.6 (20)
C1—N1—C7—H2C7	-75.8	C5—C6—C7—N1	-10.2 (17)
C2—N1—C7—C6	-79.1 (15)	C5—C6—C7—H1C7	-130.7
C2—N1—C7—H1C7	41.3	C5—C6—C7—H2C7	110.2
C2—N1—C7—H2C7	160.4	C15—C6—C7—N1	174.2 (13)
C4—N2—C1—N1	-69.8 (13)	C15—C6—C7—H1C7	53.8
C4—N2—C1—H1C1	50.5	C15—C6—C7—H2C7	-65.3
C4—N2—C1—H2C1	169.9	C5—C6—C15—C14	3.9 (21)
C5—N2—C1—N1	55.3 (16)	C5—C6—C15—HC15	-176.1
C5—N2—C1—H1C1	175.6	C7—C6—C15—C14	179.5 (12)
C5—N2—C1—H2C1	-65.0	C7—C6—C15—HC15	-0.5
C1—N2—C4—C3	42.4 (15)	Cl1—C8—C9—C10	-178.1 (12)
C1—N2—C4—H1C4	163.0	Cl1—C8—C9—HC9	1.9
C1—N2—C4—H2C4	-78.2	C2—C8—C9—C10	1.4 (23)
C5—N2—C4—C3	-81.5 (16)	C2—C8—C9—HC9	-178.6
C5—N2—C4—H1C4	39.1	C8—C9—C10—Br1	176.8 (11)
C5—N2—C4—H2C4	157.9	C8—C9—C10—C11	-3.7 (25)
C1—N2—C5—C6	-17.3 (19)	HC9—C9—C10—Br1	-3.2
C1—N2—C5—C12	156.2 (13)	HC9—C9—C10—C11	176.3
C4—N2—C5—C6	103.2 (14)	Br1—C10—C11—C3	-178.6 (12)
C4—N2—C5—C12	-83.3 (16)	Br1—C10—C11—HC11	1.4
N1—C2—C3—C4	-5.9 (21)	C9—C10—C11—C3	1.9 (26)
N1—C2—C3—C11	173.3 (14)	C9—C10—C11—HC11	-178.1
C8—C2—C3—C4	176.4 (14)	Cl2—C12—C13—C14	-176.9 (10)
C8—C2—C3—C11	-4.4 (21)	Cl2—C12—C13—HC13	3.1



## supplementary materials

---

N1—C2—C8—C11	4.3 (19)	C5—C12—C13—C14	3.7 (19)
N1—C2—C8—C9	-175.2 (13)	C5—C12—C13—HC13	-176.3
C3—C2—C8—C11	-177.9 (11)	C12—C13—C14—Br2	173.5 (9)
C3—C2—C8—C9	2.6 (21)	C12—C13—C14—C15	-2.1 (18)
C2—C3—C4—N2	-7.5 (20)	HC13—C13—C14—Br2	-6.5
C2—C3—C4—H1C4	-128.1	HC13—C13—C14—C15	177.9
C2—C3—C4—H2C4	113.1	Br2—C14—C15—C6	-177.3 (11)
C11—C3—C4—N2	173.3 (14)	Br2—C14—C15—HC15	2.7
C11—C3—C4—H1C4	52.7	C13—C14—C15—C6	-1.7 (20)
C11—C3—C4—H2C4	-66.1	C13—C14—C15—HC15	178.3

Fig. 1




Fig. 2

