

BMJ Open Forgone care among middle aged and elderly with chronic diseases in China: evidence from the China Health and Retirement Longitudinal Study Baseline Survey

Xiangjun Li,¹ Mingsheng Chen,^{2,3} Zhonghua Wang,^{2,3} Lei Si⁴

To cite: Li X, Chen M, Wang Z, *et al.* Forgone care among middle aged and elderly with chronic diseases in China: evidence from the China Health and Retirement Longitudinal Study Baseline Survey. *BMJ Open* 2018;**8**:e019901. doi:10.1136/bmjopen-2017-019901

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2017-019901>).

Received 5 October 2017
Revised 31 January 2018
Accepted 5 February 2018



¹School of Health Economics and Management, Nanjing University of Chinese Medicine, Nanjing, China

²School of Health Policy & Management, Nanjing Medical University, Nanjing, China

³Nanjing Medical University, Institute of Healthy Jiangsu Construction & Development, Nanjing, China

⁴Centre for the Health Economy, Macquarie University, New South Wales, Australia

Correspondence to
Dr Zhonghua Wang;
wzh04@njmu.edu.cn

ABSTRACT

Objective In general, published studies analyse healthcare utilisation, rather than foregone care, among different population groups. The assessment of foregone care as an aspect of healthcare system performance is important because it indicates the gap between perceived need and actual utilisation of healthcare services. This study focused on a specific vulnerable group, middle-aged and elderly people with chronic diseases, and evaluated the prevalence of foregone care and associated factors among this population in China.

Methods Data were obtained from a nationally representative household survey of middle-aged and elderly individuals (≥ 45 years), the China Health and Retirement Longitudinal Study, which was conducted by the National School of Development of Peking University in 2013. Descriptive statistics were used to analyse sample characteristics and the prevalence of foregone care. Andersen's healthcare utilisation and binary logistic models were used to evaluate the determinants of foregone care among middle-aged and elderly individuals with chronic diseases.

Results The prevalence of foregone outpatient and inpatient care among middle-aged and elderly people was 10.21% and 6.84%, respectively, whereas the prevalence of foregone care for physical examinations was relatively high (57.88%). Predisposing factors, including age, marital status, employment, education and family size, significantly affected foregone care in this population. Regarding enabling factors, individuals in the highest income group reported less foregone inpatient care or physical examinations compared with those in the lowest income group. Social healthcare insurance could significantly reduce foregone care in outpatient and inpatient situations; however, these schemes (except for urban employee medical insurance) did not appear to have a significant impact on foregone care involving physical examinations.

Conclusion In China, policy-makers may need to further adjust healthcare policies, such as health insurance schemes, and improve the hierarchical medical system, to promote reduction in foregone care and effective utilisation of health services.

Strengths and limitations of this study

- The China Health and Retirement Longitudinal Study is the first national survey focusing on middle-aged and elderly individuals, and provides detailed information about respondents and their living spouses.
- This was the first analysis of foregone care among Chinese middle-aged and elderly people with chronic diseases.
- This paper evaluated foregone care only among middle-aged and elderly adults who perceived a need for healthcare but did not seek treatment; however, it did not take into account those not perceiving the need for healthcare; therefore, the prevalence of foregone care may have been underestimated to some extent.
- Cross-sectional data cannot be used to draw causal conclusions.

INTRODUCTION

Due to the rapidly ageing population and increasing longevity, in addition to increasing improvement in medical care, a growing number of people are living for long periods of time with chronic disease. The WHO reported that chronic diseases kill 40 million people per year, which is equivalent to 70% of all deaths globally. Each year, 15 million people die from a chronic disease, and >80% of these 'premature' deaths occur in low-income and middle-income countries.¹ In recent years, the number of individuals with chronic diseases in China has continued to rise, with many residents facing serious health challenges. There were 260 million people with chronic diseases in China in 2012, and this number has increased by an average of 8.9% each year. Deaths from chronic diseases account for 86.6% of total mortality. As a result of the rapidly ageing population, the number of elderly people in China in 2050 is predicted



to total >400 million, accounting for one-third of the total population, and the average number of chronic diseases that individuals >60 years have is estimated at 2.2. China has entered a period of high economic burden caused by chronic diseases, which account for approximately 70% of the total economic disease burden.² According to a report by WHO in 2015, the direct medical costs for chronic diseases in China were >US\$500 billion, and by 2030, rapid ageing may increase the financial burden of chronic diseases by 40%.

Access to equitable healthcare, according to need, and regardless of demographic, ability to pay or social background, is an important goal for worldwide healthcare service systems.³ China has implemented a series of healthcare reforms during the past two decades, including the expansion of health insurance coverage and the reformation of the health service system, with the ultimate goal of affordable and equitable healthcare for all. The reforms of the healthcare systems have minimal benefits at the beginning and progress with expanded benefits.⁴ However, healthcare in China remains highly unequal across different subpopulations, and both the demand for, and inequality in, healthcare utilisation have increased significantly in this country, due to its rapidly ageing population. Several studies have addressed this issue by examining factors associated with healthcare utilisation among the elderly in China.^{5–8} Foregone care is another aspect of healthcare access; an individual with forgone care needs is defined as one who does not use healthcare, despite perceiving a need for it. Foregone care is an important aspect in the assessment of healthcare system performance, as it represents the gap between the perceived need and actual utilisation of healthcare services from the citizens' perspective.

Ageing is accompanied by an increase in the prevalence of non-communicable chronic diseases.⁹ In China, the elderly people are more sensitive to the economic burden of diseases overall because of the high prevalence of chronic diseases and low incomes of this population, and the absence of social security mechanisms.¹⁰ Thus, elderly individuals with chronic diseases may have a higher prevalence of forgone care compared with those who do not have such conditions. Several studies have empirically tested the associations between individual and community factors and forgone care. These studies indicate an association between higher rates of forgone care and female sex, younger age, rural living, lack of health insurance, lack of financial support, low levels of education and poor health.^{11–16} Although forgone care is an area of high interest for researchers and policy-makers, there is a dearth of research on this subject among middle-aged and elderly individuals with chronic disease in China. The overall goal of this study was to identify the prevalence of forgone care and its associated factors among middle-aged and elderly people with chronic conditions. Two specific objectives were defined: (1) comparison of the prevalence of forgone care between middle-aged and elderly individuals with and without chronic diseases

and (2) using Anderson's behavioural model, to identify factors associated with foregone care among middle-aged and elderly individuals with chronic diseases.

METHODS AND DATA

Data source/sample

This study was based on data from a nationally representative household survey of middle-aged and elderly people, the China Health and Retirement Longitudinal Study (CHARLS; <http://charls.ccer.edu.cn/en/page/data/2013-charlswave1>). CHARLS is a biennial survey conducted by the China Centre for Economic Research at Peking University. The sample was drawn in four stages. First, county-level units (counties or urban districts) were sampled directly. These counties cover 28 of 30 provinces in mainland China, other than Tibet. Second, village and community units within county units were chosen with the help of the National Bureau of Statistics, using recently updated village-level population data. The sample used administrative villages (cun) in rural areas and neighbourhoods (shequ) in urban areas as primary sampling units (PSUs). Three PSUs were selected within each county-level unit, using probability proportional to size (PPS) sampling, to select a total of 450 PSUs. Third, household units were selected in each PSU. The sampling frame was constructed using Google Earth base maps, and a computer-assisted personal interview programme was then used to sample households and to conduct the interviews using laptops. Finally, all age-eligible sample households who were willing to participate in the survey were interviewed.

CHARLS baseline data include detailed information about respondents and their living spouses. The main questionnaire includes information on basic demographics, family, health status and functioning, healthcare and insurance, work, retirement and pension, income, expenditure and assets. In the 2013 wave, the survey sample consisted of 18 605 individuals ≥45 years old and their spouses. In this paper, we excluded spouses ≤45 years old, and the remaining 18 377 observations were included in the analysis.

Statistical analysis

Binary multivariate logistic regression was applied to estimate ORs of forgone care. The dependent variable 'forgone care' was dichotomised into 0 = 'reported utilisation of healthcare' and 1 = 'reported non-use of healthcare'. All explanatory variables were recorded as binary variables.

Variable selection

The dependent variables were three kinds of forgone care: non-use of outpatient services, non-use of inpatient services and non-attendance for physical examination. Participants were asked the following two questions: 'Have you been ill in the last month?' and 'What was the main reason for not seeking outpatient treatment?', and



the responses were used to estimate 'non-use of outpatient services' (ie, if the participant answered that he/she had been ill and did not seek medical treatment). Participants were asked, 'In the past year, did a doctor suggest that you needed inpatient care but you did not get hospitalised?' and the response was used to estimate 'non-use of inpatient services' (ie, if the participants answered that they did not attend hospital when a doctor had suggested a need for inpatient services). Participants were asked, 'Did you undergo a physical examination in the last 2 years?' and the response used to estimate 'non-use of physical examination' (ie, if the participants answered that they did not undergo a physical examination in the last 2 years).

Independent variables were selected based on Andersen's healthcare utilisation model (a behavioural model). This model has guided systematic investigations into the factors that can lead to the use of healthcare services, including predisposing, enabling and need factors.¹⁷ Predisposing factors include demographic characteristics and both social structure and health belief-related factors. In our analyses, we included sex, age and marital status as demographic characteristics and education, employment status and family size as social structure factors. Enabling factors are those that make health service resources available to an individual; they can be measured by determination of family attributes, such as income, access to health insurance or other sources of third-party payment. Enabling factors in the community in which the family lives can also affect the use of services. Community attributes include the number of healthcare facilities and personnel in a community and the region of the country, in addition to the rural-urban nature of the community. In our analyses, access to public insurance, income levels and financial transfers received and given were considered as family components, while area of residence and medical facilities were considered indicators of community attributes. Need factors were both perceived need and need diagnosed or evaluated by a healthcare professional; the former was based on self-reported health status, and the latter included chronic diseases, depression, body mass index (BMI) and limitations on activities of daily living (ADL). Variables included in the multivariate model are presented in [table 1](#).

RESULTS

Sample characteristics

The total sample consisted of 18 377 respondents ≥ 45 years old. Missing data were automatically removed during data processing. No chronic disease was reported by 28.88% of respondents, with 71.12% reporting ≥ 1 chronic disease. In CHARLS, the chronic conditions included hypertension, dyslipidaemia, diabetes, cancer, chronic lung, stomach, memory-related, and/or liver diseases, heart problems, stroke, kidney, emotional and nervous problems, arthritis or rheumatism and/or asthma. Patients with chronic diseases were divided into two categories,

including individuals with one chronic and those with multiple (≥ 2) chronic diseases in one person.

Descriptive statistics of the characteristics of the individuals with, versus without, chronic conditions are presented in [table 2](#). Chronic disease prevalence was slightly higher among middle-aged and elderly women than men. Middle-aged and elderly individuals aged 45–55 years accounted for 43.28% of those without chronic disease, which was higher than the proportion of those with one (36.05%) or multiple (26.78%) chronic diseases; however, with increasing age, the proportion of individuals with one or multiple chronic diseases exceeded that of people without these conditions, indicating that the risk of chronic disease increases with age. The proportion of people without chronic diseases, and with one or multiple chronic diseases gradually increased among middle-aged and elderly people who were divorced/single/separated or unemployed/retired, indicating that the risk of chronic disease was relatively high among this group. Middle-aged and elderly people with lower educational levels ($<$ lower secondary) and small family size (≤ 2) also had a higher risk of chronic diseases.

Regarding enabling factors, the ratios of population subgroups (total, no chronic diseases, one chronic disease and multimorbidity) with social health insurance did not differ significantly. The ¥1496–8126 income group had the highest probability of having more than two types of chronic disease, and the proportion of people without chronic disease was greater in the high-income, compared with the low-income, group. Compared with middle-aged and elderly people who did not receive support from others, people with economic support exhibited an increasing trend (67.05%) towards lack of chronic disease, 73.75% towards one chronic disease and 78.14% towards multiple chronic diseases; people who provided economic support to others exhibited the opposite trend.

Among need factors, middle-aged and elderly individuals whose self-reported health was fair or poor had an increasing trend towards having no chronic disease (low probability), one or multiple chronic diseases, and those whose self-reported health was good or excellent exhibited the opposite trend. These results indicate that the risk of chronic disease was negatively associated with health status. Depression and probable depression were also associated with a significant increasing trend towards having one or more chronic diseases. There was also an increasing trend in probability from people without chronic disease, to one chronic, and then to multiple chronic diseases among middle-aged and elderly people with BMI > 24 or with any ADL.

Prevalence of foregone care

The data presented in [table 3](#) demonstrate the prevalence of foregone care among middle-aged and elderly individuals. Overall, the prevalence of non-use but ought-to-have-used outpatient services within the last 4 weeks was 10.21%, and the prevalence of non-use of inpatient services was 6.84%. Comparisons of foregone care among

**Table 1** Dependent and independent variables

	Categories	Indicators/survey questions
Dependent variables		
Non-use of outpatient services	Reported non-use of outpatient/no reported non-use of outpatient services	Questions: Have you been ill in the last month? What was the main reason for not seeking outpatient treatment?
Non-use of inpatient services	Reported non-use of inpatient/no reported non-use of inpatient services	Question: In the past year, did a doctor suggest that you needed inpatient care but you were not get hospitalised?
Non-use of physical examinations	Not undergoing physical examination/undergoing physical examination	Question: Did you have a physical examination in the last 2 years?
Independent variables		
Predisposing factors		
Sex	Male/female	
Age (years)	≤55/56–65/66–75/≥75	
Marital status	Not married/married	Question: What is your marital status?
Employment/retirement status	Employed/not employed/retired	Employed: respondents reported that they had engaged in any type of employment; not employed: not working in the last year or retired without pension; retired: retired with pension.
Education	Less than lower secondary/upper secondary and vocational training/tertiary	Question: What is your highest level of education?
Household numbers	≤2 persons in household/≥3 persons in household	Number of household members
Enabling factors		
Social health insurance	0, no social health insurance/1, UEMI/2, URMI/3, NCMI	Question: Are you the policy holder/primary beneficiary of any type of health insurance?
Income level	≤¥1495/¥1496–8126/¥8127–20 533/≥¥20 533	Yearly household income divided by the number of household members; first household member with a weight of 1, all following household members with a weight of 0.5.
Transfer received	Financial transfers received from others/no financial transfers received from others	The total amount of financial transfers that the respondent and spouse received in the past year.
Transfer given	Financial transfers given to others/no financial transfers given to others	The total amount of financial transfers that the respondent and spouse gave in the past year.
Area of residence	Urban/rural	
Need factors		
Self-reported health	Excellent/very good/good/fair/poor	Question: Would you say your health is excellent, very good, good, fair or poor?
Chronic	No chronic/one chronic/multiple chronic diseases	Question: Have you been diagnosed with any chronic diseases (conditions listed below; respond to each one) by a doctor?
Depression (Center for Epidemiological Survey (CESD) scores)	No depression (CESD<10)/probability of being depression (10≤CESD < 20)/depression (CESD≥20)	Ten questions of the CESD (Depression Scale)
BMI	≤18.5/18.624/≥24.1	BMI is the weight divided by the square of the height.
ADL	Without ADL/with any ADL	The ADLs proposed by Wallace and Herzog ³⁰ in their paper, including bathing, dressing and eating. Getting in/out of bed was added in CHARLS.
Drinking	No/yes	Whether the respondent drank alcoholic beverages in the past
Smoking	No/yes	Whether the respondent reports ever having smoked or presently smoking

ADL, activities of daily living; BMI, body mass index; CHARLS, China Health and Retirement Longitudinal Study; NCMI, new cooperative medical insurance; UEMI, urban employee medical insurance; URMI, urban resident medical insurance.

different groups demonstrated that the underutilisation rate was higher among people with multiple chronic diseases. Furthermore, the results demonstrate that individuals without chronic disease had the highest probability of not having undergone a physical examination in the last 2 years.

Multivariate regression analysis

Binary logistic models were constructed to examine the determinants of foregone care among middle-aged and elderly Chinese people with chronic diseases. Estimated coefficients describing the predicted effects of the variables are presented in [table 4](#). We also calculated overall significance values for a multivariate logistic regression

**Table 2** Sample characteristics of middle-aged and elderly people

	Total, n=18377	No chronic disease, n=5308 (28.88%)	One chronic, n=5279 (28.73%)	Multimorbidity, n=7790 (42.39%)	P values
Predisposing factors					
Sex					
Male	47.73%	50.58%	48.74%	45.11%	0.00
Female	52.27%	49.42%	51.26%	54.89%	
Age (years)					
45–55	33.76%	43.28%	36.05%	26.78%	0.00
56–65	37.85%	35.31%	37.43%	39.60%	
66–75	19.75%	13.62%	18.85%	23.88%	
≥75	8.63%	7.79%	7.66%	9.75%	
Marital status					
Divorced/single/separated	11.51%	9.07%	10.93%	13.56%	0.00
Married/partnered	88.49%	90.93%	89.07%	86.44%	
Employment/retirement status					
Employed	65.53%	75.76%	68.75%	56.57%	0.00
Unemployed	1.72%	1.58%	1.61%	1.89%	
Retired without pension	11.28%	10.18%	10.29%	12.67%	
Retired with pension	21.48%	12.48%	19.34%	28.88%	
Education					
Less than primary school	44.03%	39.43%	44.34%	46.93%	0.00
Less than lower secondary	43.07%	45.36%	43.53%	41.19%	
Upper secondary and vocational training	12.91%	15.21%	12.13%	11.87%	
Tertiary					
Family size					
≤2	34.84%	31.99%	33.12%	37.95%	0.00
≥3	65.16%	68.01%	66.88%	62.05%	
Enabling factors					
Public insurance					
No public insurance	4.70%	6.37%	4.44%	3.74%	0.00
UEMI	14.02%	13.40%	12.57%	15.43%	
URMI	7.28%	7.20%	6.64%	7.76%	
NCMI	74.00%	73.03%	76.34%	73.07%	
Income level					
≤¥1495	24.97%	24.10%	24.82%	25.56%	0.00
¥1496–8126	24.99%	22.75%	24.82%	26.38%	
¥8127–20 533	24.99%	26.36%	24.47%	24.54%	
≥¥20533	25.06%	26.80%	25.89%	23.52%	
Transfer received					
No	26.26%	32.95%	26.22%	21.86%	0.00
Yes	73.74%	67.05%	73.78%	78.14%	
Transfer given					
No	29.51%	27.46%	29.11%	31.12%	0.00
Yes	70.49%	72.54%	70.89%	68.88%	
Living area					

Continued

Table 2 Continued

	Total, n=18377	No chronic disease, n=5308 (28.88%)	One chronic, n=5279 (28.73%)	Multimorbidity, n=7790 (42.39%)	P values
Urban	37.75%	37.43%	35.85%	39.16%	0.00
Rural	62.25%	62.57%	64.15%	60.84%	
Need factors					
Self-reported of health					
Excellent	1.37%	2.97%	1.27%	0.38%	0.00
Very good	10.06%	19.23%	10.04%	3.90%	
Good	14.26%	21.82%	15.20%	8.54%	
Fair	52.51%	48.99%	56.82%	51.95%	
Poor	21.80%	6.99%	16.67%	35.24%	
Depression (CESD scores)					
No depression	68.52%	80.10%	72.24%	58.27%	0.00
Probability of being depressed	26.66%	17.93%	24.21%	34.16%	
Depression	4.82%	1.98%	3.55%	7.58%	
BMI					
≤18.5	48.74%	53.75%	51.78%	43.49%	0.00
18.6–24	5.77%	6.01%	5.87%	5.56%	
≥24.1	45.49%	40.24%	42.35%	50.96%	
ADL					
Without ADL	88.37%	92.67%	91.87%	83.06%	0.00
With any ADL	11.63%	7.33%	8.13%	16.94%	
Drinking					
Never	55.04%	53.69%	55.38%	55.72%	0.06
Ever	44.96%	46.31%	44.62%	44.28%	
Smoking					
Never	57.66%	57.03%	58.12%	57.78%	0.50
Ever	42.34%	42.97%	41.88%	42.22%	

Calculations were weighted using individual sampling weights and adjusted for household and individual responses.

ADL, activities of daily living; BMI, body mass index; CESD, Center for Epidemiological Survey, Depression; NCMI, new cooperative medical insurance; UEMI, urban employee medical insurance; URMI, urban resident medical insurance.

model (Wald χ^2 value and p value; [table 4](#)), which indicated that the overall significance of the model was strong.

With the exception of sex, all other predisposing factors were predicted to have significant impacts on foregone healthcare. Specifically, middle-aged and elderly people >55 years old were less likely to report outpatient or physical examination foregone care compared with those <55 years old. Individuals who were married or had

a larger family (>2 members) had a higher likelihood of foregone physical examination, while those with high levels of education had a significantly lower likelihood of foregone physical examination. Middle-aged and elderly people with chronic diseases who were retired without pensions exhibited a higher likelihood of outpatient and inpatient foregone care compared with retired people with pensions.

Table 3 Prevalence of foregone care among middle-aged and elderly people

Foregone care	No chronic disease	One chronic disease	Multiple chronic diseases	Total	P values
Non-use of outpatient services	362 (6.88)	482 (9.16)	1022 (13.18)	1866 (10.21)	0.0000
Non-use of inpatient services	82 (2.15)	176 (4.38)	707 (11.27)	965 (6.84)	0.0000
Non-attendance for physical examination	3372 (65.51)	3089 (60.18)	3875 (51.14)	10336 (57.88)	0.0000

Calculations were weighted using individual sampling weights and adjusted for household and individual responses.

**Table 4** Logistic regression model of forgone care among middle-aged and elderly individuals with chronic diseases

	Non-use of outpatient services		Non-use of inpatient services		Non-attendance for physical examination	
	OR	SD	OR	SD	OR	SD
Predisposing factors						
Sex, ref: female	1.06	0.16	0.94	0.21	1.04	0.11
Age (years), ref: ≤55						
56–65	0.65***	0.08	1.19	0.19	0.79***	0.07
66–75	0.56***	0.09	0.63**	0.14	0.52***	0.06
≥75	0.59*	0.17	0.66	0.24	0.52***	0.09
Marital status, ref: divorced/single/separated	0.95	0.16	0.84	0.16	1.24**	0.13
Employment/retirement status, refer: retired with pension						
Employed	1.05	0.15	0.68**	0.12	1.13	0.11
Unemployed	1.92*	0.71	1.15	0.57	1.39	0.41
Retired without pension	1.03***	0.14	1.08*	0.15	1.22	0.16
Education, ref: under primary school						
Junior school	1.11	0.13	0.1	0.15	0.92	0.07
High school and above	1.01	0.20	0.61*	0.18	0.79*	0.11
Family size, ref: ≤2	0.96	0.10	0.93	0.13	1.34***	0.09
Enabling factors						
Social health insurance, ref: no insurance						
UEMI	0.73***	0.12	0.51**	0.15	0.41***	0.09
URMI	0.81*	0.18	0.93**	0.13	0.93	0.20
NCMI	0.68**	0.15	0.58**	0.15	0.76	0.14
Income level, ref: ≤¥1495						
¥1496–8126	1.22	0.16	1.09	0.19	0.93	0.08
¥8127–20 533	1.03	0.15	0.88	0.17	0.85*	0.08
≥¥20533	0.80	0.14	0.63**	0.15	0.62***	0.07
Financial transfer received, ref: not received	1.01*	0.13	1.35	0.26	1.10	0.10
Financial transfer given, ref: not given	1.07	0.13	1.19	0.19	0.85**	0.07
Area of residence, refer: urban	1.17***	0.07	0.85	0.12	1.44***	0.12
Need factors						
Self-reported health, ref: poor						
Fair	0.85*	0.09	0.51***	0.07	1.19**	0.09
Good	0.71**	0.13	0.33***	0.09	1.21	0.14
Very good	0.52*	0.17	0.27***	0.12	0.82	0.13
Excellent	0.33	0.35	0.83	0.85	1.86	0.94
Multiple chronic, ref: one chronic	1.32***	0.14	2.74***	0.46	0.73***	0.05
Depression, ref: no depression						
Probability of being depressed	1.54***	0.17	1.42**	0.21	1.21**	0.09
Depression	1.49**	0.29	2.32***	0.55	1.18	0.17
BMI, ref: 18.6–24						
≤18.5	0.84	0.15	0.7	0.16	1.39**	0.20
≥24.1	0.60***	0.11	0.57**	0.13	1.17	0.17
ADL, ref: without ADL						
Drinking, ref: never drank	0.97	0.11	0.92	0.14	0.89	0.07
Smoking, ref: never smoked	0.95	0.14	0.95	0.20	1.23**	0.12

Continued



Table 4 Continued

	Non-use of outpatient services		Non-use of inpatient services		Non-attendance for physical examination	
	OR	SD	OR	SD	OR	SD
Wald $\chi^2(32)$	1732.42		1357.85		327.39	
Probability $>\chi^2$	0		0		0	
Observations	4928		3821		4826	

Estimates were weighted using individual sampling weights and adjusted for household and individual responses.

*P<0.05.

**P<0.01.

***P<0.001.

ADL, activities of daily living; BMI, body mass index; NCMI, new cooperative medical insurance; UEMI, urban employee medical insurance; URMI, urban resident medical insurance.

Among enabling factors, we found a significant association between social health insurance and non-use of outpatient and inpatient services; hence, enrolment in social health insurance schemes may significantly reduce the likelihood of outpatient and inpatient foregone care among middle-aged and elderly people with chronic diseases. The highest income group (\geq ¥20 533) demonstrated a significantly reduced likelihood of forgone inpatient and physical examination care compared with the lowest income group. Receipt of financial transfers increased the likelihood of non-use of outpatient services, and providing transfers was associated with reduced probability of foregone physical examination. Compared with middle-aged and elderly people living in urban areas, those in rural areas were significantly more likely to forego outpatient and physical examination care.

Among need factors, significant associations were detected between self-reported health, multiple chronic diseases, depression, and outpatient, inpatient, and physical examination foregone care. Middle-aged and elderly people with chronic diseases with fair, good or very good self-reported health had a lower likelihood of outpatient and inpatient foregone care compared with those with poor health status, and the likelihood decreased with increasing health status; however, individuals who self-reported their health as fair had a higher likelihood of foregone physical examination. Middle-aged and elderly people who had multiple chronic diseases demonstrated a higher probability of foregone outpatient and inpatient care, while exhibiting a lower likelihood of foregone physical examination. Individuals with depression had a higher likelihood of outpatient, inpatient and physical examination foregone care. Middle-aged and elderly people with BMI values >24 had a lower probability of outpatient and inpatient foregone care. Any ADL increased the possibility of outpatient service foregone care. Drinking alcohol did not have a significant impact on any kind of foregone care, while smoking significantly increased the likelihood of foregone physical examination.

DISCUSSION

Along with the rapidly ageing population in China, the number of chronic diseases continues to rise and the elderly face serious health challenges. Although many studies have analysed healthcare utilisation in diverse population groups, few have examined foregone care in middle-aged and elderly chronically ill Chinese patients. This study, focusing on middle-aged and elderly people, was performed to analyse the prevalence of foregone healthcare and its associated factors in China.

In this study, we found that 10.21% and 6.84% of respondents had foregone outpatient and inpatient care, respectively; however, the prevalence of physical examination foregone care was relatively high (57.88%). This may be attributable to the fact that Chinese social healthcare insurance schemes do not cover physical examinations; therefore, many middle-aged and elderly people are apt to forego physical examinations because of the associated economic burden. The prevalence of outpatient and inpatient foregone care rose, alongside increasing number of chronic diseases. Furthermore, the results of regression analysis (table 4) confirmed that respondents with multiple chronic diseases were more likely to forego outpatient and inpatient care. These data indicate that a greater demand for health services is associated with a higher probability of foregone outpatient and inpatient care. However, the likelihood of physical examination foregone care exhibited the opposite trend; people without chronic disease were most likely to forego physical examination and the prevalence decreased with increased number of chronic diseases. This result suggests that middle-aged and elderly individuals with chronic disease may have stronger awareness of preventative healthcare.

Among predisposing factors, our results indicate that the possibility of foregone outpatient and physical examination care decreases with increasing age. This is consistent with previous results literature reports.^{11 18} People in older age groups generally have more need for healthcare, related to more serious conditions, and may be more flexible with their time when seeking healthcare compared



with younger individuals. Middle-aged and elderly people with chronic diseases who have fixed incomes (such as old age pensions) accessed more healthcare and exhibited a relatively low likelihood of foregoing care. This finding indicates that some medical financial assistance programmes may need to be extended, particularly with respect to unemployed and elderly retired people without pensions, to facilitate further reduction of the possibility of foregone care. In addition, our analyses revealed that middle-aged and elderly people with chronic diseases who were married and had large families exhibited a relatively high probability of foregone physical examination, consistent with a number of existing literature reports.¹⁹ We did not identify any association between sex and foregone care, which is also in line with several previous publications.^{12 20}

Of enabling factors, our results demonstrate a strong association between socioeconomic variation and foregone care. Specifically, we found a significant association between household income level and foregone care; people with chronic diseases in the highest income group (\geq ¥20 533) reported less foregone inpatient and physical examination care compared with those in the lowest income group; however, we did not find a significant association between income level and outpatient foregone care, similar to previous studies.^{11 16 18} These findings indicate that due to higher hospitalisation costs, income levels remain an important impact factor influencing the possibility of inpatient foregone care; however, for outpatient services, there were no significant differences in the likelihood of foregone care among different income groups, indicating that the universal coverage of medical insurance in China may be gradually improving the equity of outpatient care utilisation among middle-aged and elderly people with chronic disease from different income levels. We also found that although social healthcare insurance can significantly reduce the level of outpatient and inpatient foregone care among chronically ill elderly patients, these schemes (except for urban employee medical insurance) may not significantly impact physical examination foregone care among elderly patients with chronic disease. Hence, policy efforts may need to focus on chronic elderly patients having urban resident medical insurance or new cooperative medical insurance, and strategies may need to be adjusted to promote the effective utilisation of preventive healthcare. Middle-aged and elderly people with chronic diseases who have received financial transfers from others exhibited a slightly higher likelihood of outpatient foregone care, while those who had made transfers to others had a significantly lower level of physical examination foregone care. Residents in rural areas had a significantly higher likelihood of outpatient and physical examination foregone care, as previously described.²¹ These findings may reflect the rural–urban income gap and differences in medical security in China.

As Andersen described, need factors best describe the true needs for healthcare.¹⁷ Consistent with previous

studies, we found that middle-aged and elderly people with multiple chronic diseases or depression were more likely to forego outpatient and inpatient care.^{8 22 23} This may be because of high demands for medical services or lack of patient compliance among this group, likely leading to a relatively high probability of foregone care. Regarding physical examinations, multiple chronic diseases were associated with a reduced possibility of foregone care; however, depression was associated with an increased likelihood, consistent with published data.^{8 24–27} Self-reported health was negatively associated with outpatient and inpatient foregone care, except for individuals whose self-reported health was excellent. This indicates that the superior health of respondents was associated with a lower likelihood of foregone care.²⁸ ADL limitations did not affect inpatient and physical examination foregone care; however, these factors significantly increased the likelihood of outpatient foregone care. This finding maybe related to the fact that people with ADL often have some difficulty seeking outpatient treatment in a timely manner.²⁹

The findings of this study may provide evidence on healthcare underutilisation and associated factors among middle-aged and elderly people. Some limitations of our study must be acknowledged. We evaluated foregone care only among middle-aged and elderly adults who perceived a need for healthcare but did not seek treatment; however, we did not take into account those who did not perceive the need for healthcare. Some patients objectively require healthcare, but do not perceive that need; therefore, the prevalence of foregone care may have been underestimated to some extent. In addition, cross-sectional data cannot be used to draw any causal conclusions about why these individuals may have foregone care.

CONCLUSION

Our findings reveal that the prevalence of foregone physical examination is relatively high among middle-aged and elderly people, and that this prevalence decreases with an increased number of chronic diseases. As a result of their greater demand for health services, respondents with chronic diseases demonstrated relatively higher likelihoods of both outpatient and inpatient foregone care. Hence, the adjustment of health insurance policies and improvements in the hierarchical medical system in China may be helpful to the effective utilisation of preventive care and outpatient and inpatient services. Some predisposing factors, including age, marital status, employment, education and family size; enabling factors, including social health insurance, income level and area of residence; and need factors, such as self-reported health, multiple chronic diseases and ADL, significantly affected foregone care among middle-aged and elderly people. Therefore, we may need to address these socioeconomic and demographic factors for reducing foregone care and enhancing the effective use of health services among

middle-aged and elderly people in China. Finally, further work, including longitudinal studies, is required to gain a better understanding of the causal relationships between foregone care and its associated factors.

Acknowledgements The authors thank the National School of Development, Peking University and other members for their support and cooperation. The authors also thank Junlong Shen and Fang Pan who provided invaluable comments and suggestions for this paper.

Contributors XL was involved in data collection, analysis and writing the manuscript. ZW was involved in data analysis and took the lead in writing the manuscript. MC analysed the data and helped to write the final draft of the manuscript. LS supervised the study, contributed to the study design and wrote the manuscript. All the authors have read and approved the entire manuscript.

Funding This study was funded by the Education Department of Jiangsu Province (Grant number: 2014SJB175).

Disclaimer The funding body was not involved in the design of the study, data collection, analysis and interpretation, or in writing the manuscript.

Competing interests None declared.

Patient consent Obtained.

Ethics approval The study was approved by the ethics committee of Nanjing Medical University; reference number: 2017613.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Extra data can be accessed via the Dryad data repository at <http://datadryad.org/> with the doi: 10.5061/dryad.n4b49kq

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Alwan A. *Global status report on non-communicable diseases 2010*. Geneva, Switzerland: World Health Organization, 2011.
- National Health and Family Planning Commission of the People's Republic of China. *China report on Nutrition and noncommunicable chronic diseases 2015*. China, Beijing: People health Press, 2015.
- WHO. The world health report. *Health systems financing: the path to universal coverage*. Geneva: World Health Organization, 2010.
- Yip WC, Hsiao WC, Chen W, *et al*. Early appraisal of China's huge and complex health-care reforms. *Lancet* 2012;379:833–42.
- Luo J, Zhang X, Jin C, *et al*. Inequality of access to health care among the urban elderly in northwestern China. *Health Policy* 2009;93:111–7.
- Niu T, Meng Q, Wang G, *et al*. Study on the health service utilization among elderly and its influencing factors in Shandong province. *Chinese Primary Health Care* 2010;24:7–8. In Chinese.
- Zhang C, Lei X, Strauss J, *et al*. Health Insurance and Health Care among the Mid-Aged and Older Chinese: Evidence from the National Baseline Survey of CHARLS. *Health Econ* 2017;26:431–49.
- Gong CH, Kendig H, He X. Factors predicting health services use among older people in China: An analysis of the China Health and Retirement Longitudinal Study 2013. *BMC Health Serv Res* 2016;16:63.
- Albanese E, Liu Z, Acosta D, *et al*. Equity in the delivery of community healthcare to older people: findings from 10/66 Dementia Research Group cross-sectional surveys in Latin America, China, India and Nigeria. *BMC Health Serv Res* 2011;11:153.
- Prince MJ, Wu F, Guo Y, *et al*. The burden of disease in older people and implications for health policy and practice. *Lancet* 2015;385:549–62.
- Röttger J, Blümel M, Köppen J, *et al*. Forgone care among chronically ill patients in Germany—Results from a cross-sectional survey with 15,565 individuals. *Health Policy* 2016;120:170–8.
- Zhou C, Ji C, Chu J, *et al*. Non-use of health care service among empty-nest elderly in Shandong, China: a cross-sectional study. *BMC Health Serv Res* 2015;15:294.
- Åhs A, Burell G, Westerling R. Care or not care—that is the question: predictors of healthcare utilisation in relation to employment status. *Int J Behav Med* 2012;19:29–38.
- Bodenmann P, Favrat B, Wolff H, *et al*. Screening primary-care patients forgoing health care for economic reasons. *PLoS One* 2014;9:e94006.
- Bryant T, Leaver C, Dunn J. Unmet healthcare need, gender, and health inequalities in Canada. *Health Policy* 2009;91:24–32.
- Ronksley PE, Sanmartin C, Quan H, *et al*. Association between chronic conditions and perceived unmet health care needs. *Open Med* 2012;6:e48–58.
- Andersen R, Newman JF. Societal and Individual Determinants of Medical Care Utilization in the United States. *Milbank Q* 2005;83:Online-only.
- Levesque JF, Pineault R, Hamel M, *et al*. Emerging organisational models of primary healthcare and unmet needs for care: insights from a population-based survey in Quebec province. *BMC Fam Pract* 2012;13:66.
- Mielck A, Kiess R, von dem Knesebeck O, *et al*. Association between forgone care and household income among the elderly in five Western European countries - analyses based on survey data from the SHARE-study. *BMC Health Serv Res* 2009;9:52.
- Hopman P, Heins MJ, Korevaar JC, *et al*. Health care utilization of patients with multiple chronic diseases in the Netherlands: Differences and underlying factors. *Eur J Intern Med* 2016;35:44–50.
- Yu H. Universal health insurance coverage for 1.3 billion people: What accounts for China's success? *Health Policy* 2015;119:1145–52.
- Peytremann-Bridevaux I, Chevrou-Severac H. Economic grand rounds: Financial burden of medical care and risk of forgoing care among Europeans with depressive symptoms. *Psychiatr Serv* 2008;59:840–2.
- Himelhoch S, Weller WE, Wu AW, *et al*. Chronic medical illness, depression, and use of acute medical services among Medicare beneficiaries. *Med Care* 2004;42:512–21.
- Srivastava D, McGuire A. Patient access to health care and medicines across low-income countries. *Soc Sci Med* 2015;133:21–7.
- Hansen AH, Halvorsen PA, Ringberg U, *et al*. Socio-economic inequalities in health care utilisation in Norway: a population based cross-sectional survey. *BMC Health Serv Res* 2012;12:336.
- Verhagen I, Steunenberg B, de Wit NJ, *et al*. Community health worker interventions to improve access to health care services for older adults from ethnic minorities: a systematic review. *BMC Health Serv Res* 2014;14:497.
- Gotsadze G, Murphy A, Shengelia N, *et al*. Healthcare utilization and expenditures for chronic and acute conditions in Georgia: does benefit package design matter? *BMC Health Serv Res* 2015;15:88.
- Bähler C, Huber CA, Brünger B, Caroline B, Carola AH, Beat B, *et al*. Multimorbidity, health care utilization and costs in an elderly community-dwelling population: a claims data based observational study. *BMC Health Serv Res* 2015;15:23.
- Jang SN, Rhee S, Cho SI. Functional Limitation and Medical Care Utilization among the Community-dwelling Elderly. *J Korean Geriatr Soc* 2007;11:198–204.
- Wallace RB, Herzog AR. Overview of the Health Measures in the Health and Retirement Study. *J Hum Resour* 1995;30:S84–S157.

BMJ Open

Forgone care among middle aged and elderly with chronic diseases in China: evidence from the China Health and Retirement Longitudinal Study Baseline Survey

Xiangjun Li, Mingsheng Chen, Zhonghua Wang and Lei Si

BMJ Open 2018 8:

doi: [10.1136/bmjopen-2017-019901](https://doi.org/10.1136/bmjopen-2017-019901)

Updated information and services can be found at:
<http://bmjopen.bmj.com/content/8/3/e019901>

These include:

References

This article cites 27 articles, 0 of which you can access for free at:
<http://bmjopen.bmj.com/content/8/3/e019901#ref-list-1>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Health economics](#) (393)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>