

Viewpoint

Eco-engineering and management strategies for marine infrastructure to reduce establishment and dispersal of non-indigenous species

Katherine A. Dafforn^{1,2}¹University of New South Wales, School of Biological, Earth and Environmental Sciences, Sydney, NSW 2052, Australia²Sydney Institute of Marine Science, Mosman, NSW 2088, AustraliaE-mail: k.dafforn@unsw.edu.au

Received: 3 February 2017 / Accepted: 3 April 2017 / Published online: 24 April 2017

Handling editor: Joana Dias

Editor's note:

This study was first presented at the 9th International Conference on Marine Bioinvasions held in Sydney, Australia, January 19–21, 2016 (<http://www.marinebioinvasions.info/previous-conferences>). Since their inception in 1999, ICMB series have provided a venue for the exchange of information on various aspects of biological invasions in marine ecosystems, including ecological research, education, management and policies tackling marine bioinvasions.

Abstract

Habitat modification and the introduction and establishment of non-indigenous species (NIS) are two of the greatest threats to global biodiversity. Human modifications of marine habitats include the introduction of boating infrastructure, coastal defences and offshore energy installations that are occurring at an increasing rate. These artificial structures are now widely recognised as providing opportunities for the establishment and dispersal of non-indigenous fouling species in new regions. This is driving increased interest into how structures might be designed and built to limit their suitability for invasive species. At the same time the potential for artificial habitats to provide habitat to native and threatened species means that the control of NIS on these structures should not just rely on antifouling. Green or eco-engineering aims to incorporate ecological theory and principles into the design of engineered structures. When combined with other management strategies that aim to increase the resistance of recipient environments there is the potential to enhance practical barriers against invaders in an increasingly developed ocean. Here I explore examples of NIS facilitation by artificial structures and the ecological theories that could be used to reduce opportunities for NIS establishment and spread. Examples include (1) manipulating the physical and chemical properties of structures to enhance native recruitment over NIS, (2) enhancing resource use of structures by native species through “pre-seeding”, (3) providing opportunities for native grazers and predators to easily access structures, and (4) considering the timing of construction/maintenance/decommissioning for artificial structures such that resources are not made available when propagule pressure is also high. These examples are not exhaustive, but rather provide a discussion point for managers of biological invasions to generate further research and application over larger spatial scales.

Key words: artificial structure, estuary, coast, offshore, community ecology, invasion theory

Introduction

Successful invasion is generally thought to comprise of three main stages; (1) a species must be transported outside of its native range (donor region) and (2) establish a self-sustaining population that is (3) able to disperse throughout a new area (recipient region) (Williamson 1996; Kolar and Lodge 2001). The introduction of non-indigenous species (NIS) is a common problem in a world increasingly connected

through infrastructure and growing transport networks (Ruiz et al. 2000; Gollasch 2002; Minchin and Gollasch 2003) and is posited as the biggest threat to biodiversity after habitat loss (Vitousek et al. 1997; Wilcove et al. 1998). As a result, there has been significant investment in reducing invasion opportunities (Hulme 2009), but where efforts to manage transport vectors or border control fail, then eradication can be costly (Pimentel et al. 2000). Hence there is a pressing need to think outside the box of conventional controls

and investigate novel solutions to reduce the invasion opportunities for NIS in our ports and harbours.

Different natural and anthropogenic selection filters act on NIS at different stages of the invasion process (Williamson and Fitter 1996; Piola et al. 2009). At the transport stage, NIS are selected for a wide environmental tolerance as they must survive entrainment and subsequent changes in environmental conditions from the donor to recipient region (Johnston et al. 2009; Keller et al. 2011). Successful establishment will then depend on them finding a “niche opportunity” within the new region (Tilman 1997; Shea and Chesson 2002). As a result only a subset of transported species is actually able to establish and persist. This has also been defined in statistical terms as the “tens” rule (Williamson et al. 1986; Williamson and Fitter 1996) and refers to the probability that only one in every ten species transported is likely to arrive at a new destination, and only one in ten of those arrivals will establish and disperse in the new region. While we have some idea of the selection agents acting on species introductions (Williamson and Fitter 1996), further investigation is needed to determine how these selection agents could be manipulated in marine developments for the management of invasions.

The increasing construction of estuarine, coastal and offshore infrastructure provides novel habitat that has been shown to increase “niche opportunity” for NIS (Bulleri and Airolidi 2005; Bulleri et al. 2006; Glasby et al. 2007; Dafforn et al. 2009b; Airolidi and Bulleri 2011; Dafforn et al. 2012; Simkanin et al. 2012; Airolidi et al. 2015). This has been extensively reviewed elsewhere (Mineur et al. 2012; Johnston et al. 2017), but briefly can occur through the provision of physical and chemical conditions that match the environmental tolerance of arriving NIS, increased availability of limiting resources and reduced resistance in the form of natural enemies. Ecological strategies are increasingly being considered by marine developers in efforts to build and marine infrastructure in a more environmentally sustainable way. Most examples where ecological theory has been used to enhance engineered structures are related to manipulations of physical properties such as material, complexity, or microhabitat addition (“green” or “eco-engineering” e.g. Chapman and Blockley 2009; Browne and Chapman 2011; Chapman and Underwood 2011; Browne and Chapman 2014; Dafforn et al. 2015a; Firth et al. 2016b; Firth et al. 2016c; Morris et al. 2016; Mayer-Pinto et al. 2017). There is also a growing number of strategies to manage marine infrastructure that could be applied to reduce NIS threats including marine spatial planning (Gilbert et al. 2015), water quality improvement (Piola et al. 2009; Johnston et

al. in press) and implementation of regulatory drivers (Naylor et al. 2012). These strategies together with interdisciplinary approaches to marine development provide significant capacity to design, build, and manage structures that not only fulfil their major objective as physical infrastructure, but also reduce “niche opportunities” for NIS (Sella and Perkol-Finkel 2015).

Below are examples where built infrastructure (or artificial structures) have supported the establishment and dispersal of marine NIS. I use community ecology theory (Shea and Chesson 2002) as a framework for identifying the properties of built infrastructure that (1) have facilitated the establishment and dispersal of NIS, and (2) could be manipulated through eco-engineering to prevent future invasions. I provide practical small-scale examples that have been or could be incorporated into the construction or retrofit of artificial structures and outline future directions for research.

Marine infrastructure facilitates the establishment and dispersal of non-indigenous species

In areas of Europe, the United States, Australia and Asia large sections of the coastline is now modified by the addition of built infrastructure such as seawalls, breakwaters, pilings and marinas (Bulleri and Chapman 2010; Dugan et al. 2011; Dafforn et al. 2015a). Together with this, global demand for resources has driven exploration and extraction offshore where aquaculture and energy platforms are creating an increasing footprint in the ocean (Wilson and Elliott 2009; Wilson et al. 2010; Firth et al. 2016a; Heery et al. in press). Duarte (2012) refers to this collectively as “ocean sprawl” because human activities once confined to land continue to expand into ocean ecosystems. Since NIS are closely associated with artificial structures, ocean sprawl has been and will continue to be a major factor in their global spread.

Marine infrastructure, including ports, marinas, breakwaters, oil and gas platforms, has been found to be highly susceptible to NIS establishment. This is reviewed extensively in Mineur et al. (2012), but some examples include boating infrastructure (pilings and pontoons), which has previously been found to support up to 2.5× more NIS than natural reef (Glasby et al. 2007), with pontoons in particular acting as invader hotspots (Dafforn et al. 2009b). Moreover, energy infrastructure (oil platforms and rigs) can support communities comprising between 20–30% NIS (Ferreira et al. 2006; Hopkins and Forrest 2010). Apart from the provision of local habitat for the recruitment of NIS, recent research

has highlighted the role of artificial structures in their spread. For example, the northward range shift of the non-indigenous coral *Oculina patagonica* along 400 km of Mediterranean coastline is thought to have occurred more rapidly on artificial than natural habitats (Serrano et al. 2013). Similarly, the non-indigenous alga *Undaria pinnatifida* has experienced a northward range expansion from its southernmost introduction point in France to Northern Ireland in just 40 years (Minchin and Nunn 2014). While the initial introduction is thought to have been with the oyster aquaculture industry, the spread of the alga through Europe has been linked to recreational craft and the European network of marinas. Airoidi et al. (2015) also provide compelling evidence that coastal infrastructure along 500 km of coastline in the North Adriatic sea has influenced the regional spread of non-indigenous ascidians. Together these examples provide a convincing argument for designing infrastructure in more ecologically-sustainable ways, specifically to limit the establishment and dispersal of NIS.

“Niche opportunities” provided by artificial structures

Following the framework proposed by Shea and Chesson (2002), artificial structures potentially facilitate invasion through a number of processes. Firstly, through the construction of habitats with physico-chemical conditions, which match the characteristics of individual or multiple NIS (habitat suitability, Moyle and Light 1996). Secondly, through the provision of space that may be limiting in the recipient region (resource availability, Davis et al. 2000), and thirdly, by reducing biotic resistance from native species and natural enemies (biotic resistance and enemy release, Keane and Crawley 2002; Levine et al. 2004). These processes also have the potential to act and interact throughout the life cycle of an artificial structure, from construction and operation to decommissioning.

The construction of infrastructure adds substrates to the marine environment that are physically and chemically different to natural habitats. Artificial structures are often oriented vertically and shaded artificially (Glasby and Connell 1999), therefore they can experience less light than natural habitats (Glasby 1999). The design of artificial structures has also been linked to changes in currents around them, predominantly reduced flow which could increase encounter probabilities for NIS and the chance of establishment (Glasby et al. 2007). These properties are especially relevant since a large majority of NIS are fouling invertebrates and benefit from reduced

algal competition under low light conditions (Dafforn et al. 2012). Lower levels of sedimentation and slower flow under these conditions also benefit invertebrates that rely on filtering apparatus to obtain food (Ostroumov 2005). Similarly NIS establishment and spread on marina structures like pontoons can be explained by their depth range and floating position in the water column (Glasby et al. 2007), which together create physical (e.g. floating) or chemical (e.g. metal contamination in the water column) conditions for marine invaders that are analogous to the boat hulls upon which they may have arrived (Dafforn et al. 2009b). The materials used in construction may also be toxic e.g. pilings treated with copper chromated arsenate (Weis and Weis 1992) or chemically inhibit biological recruitment due to high surface alkalinity e.g. Portland cement used in breakwaters (Sella and Perkol-Finkel 2015). While copper is widely accepted to facilitate NIS dominance over native species, the potential for concrete mixtures to enhance recruitment of target species on artificial structures has only recently begun to be explored (Sella and Perkol-Finkel 2015).

Artificial structures also increase the availability of a limiting resource in marine environments, i.e. space through the provision of hard substrates (Dayton 1971; Buss 1979; Bullard et al. 2004; Glasby et al. 2007). Since many fouling NIS are fast-growing or opportunistic (r-strategists), free space could be enough to facilitate their establishment over slower growing native species (k-strategists) (Grime 1977). In other instances, the provision of resources needs to also match the physico-chemical requirements of the NIS. For example, establishment of the non-indigenous alga *Codium fragile* ssp. *tomentosoides* in the North Adriatic Sea has been linked to the space provided by breakwaters, particularly on sheltered (landward) sections and with disturbances that reduced biotic resistance from dominant native mussels (Bulleri and Airoidi 2005).

During operation and decommissioning, artificial structures also change the local environmental conditions and may create conditions that favour NIS. For example, Rivero et al. (2013) found that the construction of a breakwater around a marina can reduce flow by 30%. This traps contaminants leaching from antifouling paints and has the potential to entrain the larvae and spores. Enclosed boat harbours and marinas can therefore facilitate transport of a native species to a new region, where it is non-indigenous, but also can act as a propagule source for regional dispersal (Floerl and Inglis 2003). Furthermore, boat harbours contaminated with metals from antifouling paints have been shown to facilitate invader dominance over native species

(Piola and Johnston 2008). When in combination, the presence of metal-tolerant NIS in low flow conditions at contaminated sites (e.g. marinas) can contribute to these systems being more invaded and acting as a major source of intraregional dispersal (reviewed by Johnston et al. 2017).

Hydrodynamic modelling of energy platforms in the North Sea have also shown that changes to flow as a result of the underwater scaffolding may trap propagules (larvae) that would otherwise be lost offshore, and when these propagules are non-indigenous, then potentially facilitate establishment and dispersal through oil fields (Adams et al. 2014). The trapping of propagules in areas where artificial substrate is readily available (space is not limiting) and shaded (physico-chemical conditions are suitable for fouling species) could further facilitate NIS establishment over the propagules of any native species that might also be present. Furthermore, there is a continuing argument as to whether non-operational energy platforms should be decommissioned or remain as reefs (Macreadie et al. 2011) that could potentially support NIS.

Eco-engineering and other management strategies to reduce “niche opportunities” for non-indigenous species on marine infrastructure

There are increasing opportunities for marine researchers, managers and developers to work together to reduce marine invasion through ecologically-informed design, construction and management of artificial structures (Naylor et al. 2012). At its most base level this could include regulations that control the timing of different stages of marine developments. For example, the construction of a new structure in the marine environment could consider major recruitment patterns of NIS and avoid providing an important resource at times when free space might increase the chance of successful settlement (Lockwood et al. 2005; Clark and Johnston 2009; Hedge and Johnston 2012). Given the cost and scale of many developments, timing controls at the initial construction stage might not be logistically possible. The ongoing maintenance of existing artificial structures offers more possibilities for timing control to prevent disturbances that free up resources when NIS can take advantage of them (Airoidi and Bulleri 2011). Physical cleaning practices, such as in-water cleaning, should be carefully considered in maintenance scheduling to avoid causing NIS to spawn in reproductive periods and increase local propagule pressure (Hopkins and Forrest 2008).

There is a long history of biofouling control in the marine environment and much of this has been driven by shipping, aquaculture and port industries, where the removal of all fouling species is desirable to prevent damage to important infrastructure and avoid economic costs (Dafforn et al. 2011). Strategies have included non-discriminant antifouling applications, chemical control and physical removal. While there have been demonstrated examples that these strategies can be effective controls of fouling and NIS (Champ and Pugh 1987), negative impacts (e.g. from metal pollution) on native species have also occurred (Terlizzi et al. 2001). Increasingly research suggests that manipulating substrate composition, micro- and macro-topography might offer a non-toxic alternative for directly preventing NIS recruitment while facilitating native species (Tyrrell and Byers 2007; Sella and Perkol-Finkel 2015; Strain et al. unpublished data). Furthermore, manipulations of substrate can indirectly affect the recruitment of NIS and native species in several ways, including facilitation of native species that compete with NIS for space (Sella and Perkol-Finkel 2015), and changes to the biofilm (Tan et al. 2015). The potential for microbial assemblages to be manipulated to prevent NIS recruitment is an important area of research, and molecular advances now give us an opportunity to understand how eco-engineering designs might influence fouling communities from the biofilm upwards.

Larger-scale physical manipulations of artificial structures could also be used to manage NIS recruitment and spread. For example, there is significant evidence that shading from artificial structures facilitates a predominantly invertebrate community (Miller and Etter 2008; Dafforn et al. 2012), and that NIS comprise a significant proportion of these shaded communities (Wasson et al. 2005). Therefore designs that aim to reduce shading, such as “skylights” in foreshore boardwalks composed of metal grills or transparent materials (Dyson and Yocom 2014), could go some way to encouraging native algal recruitment and NIS exclusion. Similarly, building structures such as seawalls on a more gentle gradient could create conditions more similar to natural reefs systems (Department of Environment and Climate Change 2009), reduce shading, and therefore create physico-chemical conditions that are more suitable for native than NIS establishment (Dafforn et al. 2015a).

On a larger scale, the placement of artificial structures locally, regionally and internationally should be incorporated into marine spatial planning (Dafforn et al. 2015b; Domínguez-Tejo et al. 2016). Further research is needed to e.g. determine appropriate densities, sizes and connectivity of artificial

structures that could potentially allow them to provide a pathway for climate migrants (species extending their ranges due to climate change Vergés et al. 2014), but not act as stepping stones (Airoldi et al. 2015) for the dispersal of NIS (Bishop et al. in press). Breakwaters, locks and weirs are often used to protect important infrastructure, and while they have been linked to conditions with reduced flow and propagule entrainment (Floerl and Inglis 2003; Rivero et al. 2013), there is also the potential that at ecologically-sensitive sites such as marine protected areas, they could be used to physically prevent NIS spread by trapping propagules or containing them during an eradication effort (e.g. Willan et al. 2000). An important trade-off associated with these physical interventions is the build-up of contaminants such as the metals from antifouling paints used on boats that creates degraded conditions (Sim et al. 2015).

Ports and harbours are particularly impacted by chemical contaminants (Wamken et al. 2004; Dafforn et al. 2008; Dafforn et al. 2009a). The sources of these include land-based stormwater inputs such as metals and nutrients (Birch 2000), legacy inorganic and organic contaminants from past industrial practices (Birch 2000; Jones 2000; Maguire 2000), and marine-based contaminants from antifouling paints and vessel discharges (Schiff et al. 2004). Chemical disturbance in estuaries is posited to be a major factor in the loss of native species' resistance to invasion (Piola and Johnston 2008; Johnston et al. in press) and can be beneficial for opportunistic invaders that have arrived on the hulls of vessels and already been selected for metal tolerance (Piola et al. 2009). Measures that are increasingly being incorporated into foreshore developments include water sensitive urban design that often aims to trap and recycle all contaminants through measures such as artificial wetlands and filtration systems (Hatt et al. 2006; Davis and Birch 2009; Madhani and Brown 2015; Prosser et al. 2015; Greenway in press). Copper contamination in particular has been implicated in the establishment of NIS (Piola et al. 2009). Apart from being a major contaminant in stormwater runoff, copper is also the primary component of many antifouling paints, and wooden pilings are often treated with copper chromated arsenate to prevent fouling (Weis et al. 1993). Limiting the use of copper in antifouling products would go some way to reducing chemical advantages for NIS establishment and dispersal, and helping to rebuild native species resistance to invasion (Piola et al. 2009; Johnston 2011; Johnston et al. in press).

Apart from physical and chemical manipulations, the biotic characteristics of artificial structures could be directly manipulated to increase resource use and

enhance biotic resistance, thereby decreasing opportunities for NIS. "Pre-seeding" artificial structures with native species is gathering traction in eco-engineering (Perkol-Finkel et al. 2012) and has analogous principles to the cultivation of native plant communities on green roofs or walls (Johnson and Newton 1996; Oberndorfer et al. 2007). Research has demonstrated that pre-empting space with dominant species such as canopy-forming algae can reduce invasion by hard substrate invertebrates (Dafforn et al. 2012) and also provide a refuge for threatened native species (Perkol-Finkel et al. 2012). The lack of natural enemies, such as benthic grazers and predators, on many artificial structures has also been experimentally shown to influence successful utilisation of these habitats by NIS (Dumont et al. 2011; Forrest et al. 2013; Simkanin et al. 2013). Relocating mobile species and "pre-seeding" structures with key grazers or predators appropriate to the system e.g. gastropods and/or echinoderms could go some way to enhancing biotic resistance on artificial structures. Building structures with a gentle gradient or that enhance connectivity with the benthos (Bishop et al. in press) might also provide more natural opportunities for benthic predators to extend their feeding range to artificial structures.

Below are examples of how experimental findings from invasion ecology could be applied in an eco-engineering context. These include small-scale experimental examples that demonstrate the potential for building structures with specific physical and chemical properties that directly or indirectly reduce colonisation by NIS; manipulating assemblages on structures to reduce resource availability; and improving biotic resistance of the resident community to reduce NIS establishment, while also supporting native species:

1. Building with natural substrates instead of artificial substrates can reduce space occupation by NIS up to 54% and increase native species by 45% over time (Tyrrell and Byers 2007);
2. Building with natural bedrock (e.g. sandstone in Sydney Harbour) can reduce NIS richness by ~ 30% compared to concrete or wood (Glasby et al. 2007);
3. Building with specially formulated concrete (e.g. EConcrete[®]) compared to standard concrete can reduce NIS richness by up to 50% and increase native species richness by ~ 43% (Sella and Perkol-Finkel 2015);
4. Building fixed rather than shallow, floating (e.g. pontoon) structures can reduce NIS richness by 28% (Dafforn et al. 2009b);

5. Reducing shading (e.g. by building on a gentle gradient or adding light penetrating features to over-water structures) could reduce NIS dominance by 50–90% (Kim and Micheli 2013);
6. Controlling metal contaminant inputs (e.g. stormwater management, antifouling regulations) could increase native species richness by 34% and reduce NIS dominance (space occupation) by up to 29% (Piola and Johnston 2008);
7. “Pre-seeding substrates” with native fouling species (e.g. habitat forming algae) can reduce occupation by NIS by up to 33% (Dafforn et al. 2012);
8. Increasing native predator abundance on artificial structures (e.g. by increasing connectivity to benthos) can reduce NIS dominance (e.g. ascidians) by ~ 50% (Simkanin et al. 2013).

While these examples provided above show great promise for application in marine developments, so far most have only been investigated at a small-scale and require testing over larger spatial scales. Furthermore, the logistical difficulties of each strategy to apply over a large-scale will vary with location. For example, the application of natural substrates (sandstone) in the eco-engineering design at Headland Park, Sydney Harbour required a significant rock supply to build 1.4 km of intertidal foreshore. This was cut from the headland during construction of a nearby underground car park because local sandstone supplies are limited. Where natural substrates are not an option, specially formulated concrete could provide a more readily available alternative and this has already been investigated at the scale of breakwater sections (Sella and Perkol-Finkel 2015). In comparison, biological manipulations such as “pre-seeding” are likely to prove more difficult than physical manipulations to apply over large spatial scales due to e.g. mortality during transport and the difficulty of establishment at target sites. Despite this, breakwater-scale testing has been done with a threatened algal species (Perkol-Finkel et al. 2012) and restoration projects show increasing success at large-scales (Campbell et al. 2014).

Conclusion

Evidence so far that eco-engineering could be used to control NIS success is largely experimental and small-scale. Eco-engineering will not provide a rapid solution for managing NIS establishment and dispersal. For it to become widely adopted and effective over the longer term requires a change in management principles, policy and regulation, interdisciplinary communication and co-ordination. Practical solutions for reducing niche opportunities

on artificial structures therefore need to be explored over longer timescales and larger spatial scales. Future research should investigate reducing niche opportunities at each stage of a marine development (construction, operation, decommissioning), and also at each stage of a fouling community development. The latter will benefit from (1) advances in molecular techniques that can characterise the “unseen” and unculturable diversity on these structures to improve understanding of how the early biofilm development might determine later dominance of native or NIS, and (2) new strategies to inoculate or “pre-seed” structures with relevant species and/or communities. Future population growth and resource demand will place increasing pressure on our marine environment and better integration of ecological theory and principles has the potential to mitigate invasion and provide positive benefits.

Acknowledgements

Thanks to the organisers and sponsors of the International Conference on Marine Bioinvasions (ICMB) for the opportunity to present this viewpoint and to the editors of Management of Biological Invasions for the invitation to submit this viewpoint to the Special Issue for the ICMB. I would also like to thank three anonymous reviewers for their feedback on the manuscript. This is SIMS publication # 207.

References

- Adams TP, Miller RG, Aleynik D, Burrows MT (2014) Offshore marine renewable energy devices as stepping stones across biogeographical boundaries. *Journal of Applied Ecology* 51: 330–338, <https://doi.org/10.1111/1365-2664.12207>
- Airoldi L, Bulleri F (2011) Anthropogenic disturbance can determine the magnitude of opportunistic species responses on marine urban infrastructures. *PLoS ONE* 6: e22985, <https://doi.org/10.1371/journal.pone.0022985>
- Airoldi L, Turon X, Perkol-Finkel S, Rius M (2015) Corridors for aliens but not for natives: effects of marine urban sprawl at a regional scale. *Diversity and Distributions* 21: 755–768, <https://doi.org/10.1111/ddi.12301>
- Birch GF (2000) Marine pollution in Australia, with special emphasis on central New South Wales estuaries and adjacent continental margin. *International Journal of Environment and Pollution* 13: 573–607, <https://doi.org/10.1504/IJEP.2000.002334>
- Bishop MJ, Mayer-Pinto M, Airoldi L, Firth LB, Morris RL, Loke LHL, Hawkins SJ, Naylor LA, Coleman RA, Chee SY, Dafforn KA (in press) Effects of ocean sprawl on ecological connectivity: impacts and solutions. *Journal of Experimental Marine Biology and Ecology*, <https://doi.org/10.1016/j.jembe.2017.01.021>
- Browne MA, Chapman MG (2011) Ecologically informed engineering reduces loss of intertidal biodiversity on artificial shorelines. *Environmental Science & Technology* 45: 8204–8207, <https://doi.org/10.1021/es201924b>
- Browne MA, Chapman MG (2014) Mitigating against the loss of species by adding artificial intertidal pools to existing seawalls. *Marine Ecology Progress Series* 497: 119–129, <https://doi.org/10.3354/meps10596>
- Bullard SG, Whitlatch RB, Osman RW (2004) Checking the landing zone: Do invertebrate larvae avoid settling near superior spatial competitors? *Marine Ecology Progress Series* 280: 239–247, <https://doi.org/10.3354/meps280239>

- Bulleri F, Abbiati M, Airoldi L (2006) The colonisation of human-made structures by the invasive alga *Codium fragile* ssp. *tomentosoides* in the north Adriatic Sea (NE Mediterranean). *Hydrobiologia* 555: 263–269, <https://doi.org/10.1111/j.1365-2664.2005.01096.x>
- Bulleri F, Airoldi L (2005) Artificial marine structures facilitate the spread of a non-indigenous green alga, *Codium fragile* ssp. *tomentosoides*, in the north Adriatic Sea. *Journal of Applied Ecology* 42: 1063–1072, <https://doi.org/10.1111/j.1365-2664.2005.01096.x>
- Bulleri F, Chapman MG (2010) The introduction of coastal infrastructures as a driver of change in marine environments. *Journal of Applied Ecology* 47: 26–35, <https://doi.org/10.1111/j.1365-2664.2009.01751.x>
- Buss LW (1979) Bryozoan overgrowth interactions - the interdependence of competition for space and food. *Nature* 281: 475–477, <https://doi.org/10.1038/281475a0>
- Campbell AH, Marzinielli EM, Vergés A, Coleman MA, Steinberg PD (2014) Towards restoration of missing underwater forests. *PLoS ONE* 9: e84106, <https://doi.org/10.1371/journal.pone.0084106>
- Champ MA, Pugh WL (1987) Tributyltin antifouling paints: introduction and overview. In: Proceedings of the Oceans '87 International Organotin Symposium, Washington, DC, pp 1296–1308, <https://doi.org/10.1109/oceans.1987.1160644>
- Chapman MG, Blockley DJ (2009) Engineering novel habitats on urban infrastructure to increase intertidal biodiversity. *Oecologia* 161: 625–635, <https://doi.org/10.1007/s00442-009-1393-y>
- Chapman MG, Underwood AJ (2011) Evaluation of ecological engineering of “armoured” shorelines to improve their value as habitat. *Journal of Experimental Marine Biology and Ecology* 400: 302–313, <https://doi.org/10.1016/j.jembe.2011.02.025>
- Clark GF, Johnston EL (2009) Propagule pressure and disturbance interact to overcome biotic resistance of marine invertebrate communities. *Oikos* 118: 1679–1686, <https://doi.org/10.1111/j.1600-0706.2009.17564.x>
- Dafforn KA, Glasby TM, Johnston EL (2008) Differential effects of tributyltin and copper anti-foulants on recruitment of non-indigenous species. *Biofouling* 24: 23–33, <https://doi.org/10.1080/08927010701730329>
- Dafforn KA, Glasby TM, Johnston EL (2009a) Links between estuarine condition and spatial distributions of marine invaders. *Diversity and Distributions* 15: 807–821, <https://doi.org/10.1111/j.1472-4642.2009.00587.x>
- Dafforn KA, Johnston EL, Glasby TM (2009b) Shallow moving structures promote marine invader dominance. *Biofouling* 25: 277–287, <https://doi.org/10.1080/08927010802710618>
- Dafforn KA, Lewis JA, Johnston EL (2011) Antifouling strategies: History and regulation, ecological impacts and mitigation. *Marine Pollution Bulletin* 62: 453–465, <https://doi.org/10.1016/j.marpolbul.2011.01.012>
- Dafforn KA, Glasby TM, Johnston EL (2012) Comparing the invisibility of experimental “reefs” with field observations of natural reefs and artificial structures. *PLoS ONE* 7: e38124, <https://doi.org/10.1371/journal.pone.0038124>
- Dafforn KA, Glasby TM, Airoldi L, Rivero NK, Mayer-Pinto M, Johnston EL (2015a) Marine urbanisation: an ecological framework for designing multifunctional artificial structures. *Frontiers in Ecology and the Environment* 13: 82–90, <https://doi.org/10.1890/140050>
- Dafforn KA, Mayer-Pinto M, Morris RL, Waltham NJ (2015b) Application of management tools to integrate ecological principles with the design of marine infrastructure. *Journal of Environmental Management* 158: 61–73, <https://doi.org/10.1016/j.jenvman.2015.05.001>
- Davis BS, Birch GF (2009) Catchment-wide assessment of the cost-effectiveness of stormwater remediation measures in urban areas. *Environmental Science & Policy* 12: 84–91, <https://doi.org/10.1016/j.envsci.2008.09.004>
- Davis MA, Grime JP, Thompson K (2000) Fluctuating resources in plant communities: a general theory of invasibility. *Journal of Ecology* 88: 528–534, <https://doi.org/10.1046/j.1365-2745.2000.00473.x>
- Dayton PK (1971) Competition, disturbance, and community organization: the provision and subsequent utilization of space in a rocky intertidal community. *Ecological Monographs* 41: 351–389, <https://doi.org/10.2307/1948498>
- Department of Environment and Climate Change (2009) Environmentally Friendly Seawalls: A Guide to Improving the Environmental Value of Seawalls and Seawall-lined Foreshores in Estuaries. NSW Department of Environment and Climate Change, Sydney, Australia, pp 1–34
- Dominguez-Tejo E, Metternicht G, Johnston EL, Hedge LH (2016) Marine Spatial Planning advancing the Ecosystem-Based Approach to coastal zone management: A review. *Marine Policy* 72: 115–130, <https://doi.org/10.1016/j.marpol.2016.06.023>
- Duarte CM, Pitt KA, Lucas CH, Purcell JE, Uye S-I, Robinson K, Brotz L, Decker MB, Sutherland KR, Malej A, Madin L, Mianzan H, Gili J-M, Fuentes V, Atienza D, Pagés F, Breitbart D, Malek J, Graham WM, Condon RH (2012) Is global ocean sprawl a cause of jellyfish blooms? *Frontiers in Ecology and the Environment* 11: 91–97, <https://doi.org/10.1890/110246>
- Dugan JE, Airoldi L, Chapman MG, Walker SJ, Schlacher T (2011) 8.02 - Estuarine and coastal structures: Environmental effects, a focus on shore and nearshore structures. In: Wolanski E, McLusky D (eds), Treatise on Estuarine and Coastal Science, Academic Press, Waltham, pp 17–41
- Dumont CP, Harris LG, Gaymer CF (2011) Anthropogenic structures as a spatial refuge from predation for the invasive bryozoan *Bugula neritina*. *Marine Ecology Progress Series* 427: 95–103, <https://doi.org/10.3354/meps09040>
- Dyson K, Yocom K (2014) Ecological design for urban waterfronts. *Urban Ecosystems* 18: 189–208, <https://doi.org/10.1007/s11252-014-0385-9>
- Ferreira CEL, Gonçalves JEA, Coutinho R (2006) Ship hulls and oil platforms as potential vectors to marine species introduction. *Journal of Coastal Research* 39: 1340–1345, <http://www.jstor.org/stable/25742972>
- Firth LB, Knights AM, Thompson RC, Mieszkowska N, Bridger D, Evans AJ, Moore PJ, O'Connor N, Sheehan E, Hawkins SJ (2016a) Ocean sprawl: challenges and opportunities for biodiversity management in a changing world. *Oceanography and Marine Biology Annual Review*, <https://doi.org/10.1201/9781315368597-5>
- Firth LB, Browne KA, Knights AM, Hawkins SJ, Nash R (2016b) Eco-engineered rock pools: a concrete solution to biodiversity loss and urban sprawl in the marine environment. *Environmental Research Letters* 11: 094015, <https://doi.org/10.1088/1748-9326/11/9/094015>
- Firth LB, White FJ, Schofield M, Hanley ME, Burrows MT, Thompson RC, Skov MW, Evans AJ, Moore PJ, Hawkins SJ (2016c) Facing the future: the importance of substratum features for ecological engineering of artificial habitats in the rocky intertidal. *Marine and Freshwater Research* 67: 131–143, <https://doi.org/10.1071/MF14163>
- Floerl O, Inglis GJ (2003) Boat harbour design can exacerbate hull fouling. *Austral Ecology* 28: 116–127, <https://doi.org/10.1046/j.1442-9993.2003.01254.x>
- Forrest BM, Fletcher LM, Atalah J, Piola RF, Hopkins GA (2013) Predation limits spread of *Didemnum vexillum* into natural habitats from refuges on anthropogenic structures. *PLoS ONE* 8: e82229, <https://doi.org/10.1371/journal.pone.0082229>
- Gilbert AJ, Alexander K, Sardá R, Brazinskaite R, Fischer C, Gee K, Jessopp M, Kershaw P, Los HJ, March Morla D, O'Mahony C, Pihlajamäki M, Rees S, Varjopuro R (2015) Marine spatial planning and Good Environmental Status: a perspective on spatial and temporal dimensions. *Ecology and Society* 20: 64, <https://doi.org/10.5751/es-06979-200164>
- Glasby TM (1999) Effects of shading on subtidal epibiotic assemblages. *Journal of Experimental Marine Biology and Ecology* 234: 275–290, [https://doi.org/10.1016/S0022-0981\(98\)00156-7](https://doi.org/10.1016/S0022-0981(98)00156-7)
- Glasby TM, Connell SD (1999) Urban structures as marine habitats. *Ambio* 28: 595–598

- Glasby TM, Connell SD, Holloway MG, Hewitt CL (2007) Nonindigenous biota on artificial structures: could habitat creation facilitate biological invasions? *Marine Biology* 151: 887–895, <https://doi.org/10.1007/s00227-006-0552-5>
- Gollasch S (2002) The importance of ship hull fouling as a vector of species introductions into the North Sea. *Biofouling* 18: 105–121, <https://doi.org/10.1080/08927010290011361>
- Greenway M (in press) Stormwater wetlands for the enhancement of environmental ecosystem services: case studies for two retrofit wetlands in Brisbane, Australia. *Journal of Cleaner Production*, <https://doi.org/10.1016/j.jclepro.2015.12.081>
- Heery EC, Bishop MJ, Critchley L, Bugnot AB, Airoidi L, Mayer-Pinto M, Sheehan EV, Coleman RA, Loke L, Johnston EL, Komyakova V, Morris RL, Strain E, Naylor LA, Dafforn KA (in press) Identifying the consequences of ocean sprawl for sedimentary habitats. *Journal of Experimental Marine Biology and Ecology*, <https://doi.org/10.1016/j.jembe.2017.01.020>
- Grime JP (1977) Evidence for existence of 3 primary strategies in plants and its relevance to ecological and evolutionary theory. *American Naturalist* 111: 1169–1194, <https://doi.org/10.1086/283244>
- Hatt BE, Deletic A, Fletcher TD (2006) Integrated treatment and recycling of stormwater: a review of Australian practice. *Journal of Environmental Management* 79: 102–113, <https://doi.org/10.1016/j.jenvman.2005.06.003>
- Hedge LH, Johnston EL (2012) Propagule pressure determines recruitment from a commercial shipping pier. *Biofouling* 28: 73–85, <https://doi.org/10.1080/08927014.2011.652622>
- Hopkins GA, Forrest BM (2008) Management options for vessel hull fouling: an overview of risks posed by in-water cleaning. *ICES Journal of Marine Science: Journal du Conseil* 65: 811–815, <https://doi.org/10.1093/icesjms/fsn026>
- Hopkins GA, Forrest BM (2010) Challenges associated with pre-border management of biofouling on oil rigs. *Marine Pollution Bulletin* 60: 1924–1929, <https://doi.org/10.1016/j.marpolbul.2010.07.015>
- Hulme PE (2009) Trade, transport and trouble: managing invasive species pathways in an era of globalization. *Journal of Applied Ecology* 46: 10–18, <https://doi.org/10.1111/j.1365-2664.2008.01600.x>
- Johnson J, Newton J (1996) Building Green, a guide for using plants on roofs and pavement. The London Ecology Unit, London, UK, pp 1–94
- Johnston E (2011) Tolerance to contaminants: Evidence from chronically-exposed populations of aquatic organisms. In: Amiard-Triquet C, Rainbow PS, Romeo M (eds), Tolerance to Environmental Contaminants. CRC Press, Boca Raton, pp 25–46
- Johnston EL, Piola RF, Clark GF (2009) The role of propagule pressure in invasion success. In: Rilov G, Crooks J (eds), Biological Invasions in Marine Ecosystems: Ecological, Management, and Geographic Perspectives. Springer-Verlag, Berlin, pp 133–151, https://doi.org/10.1007/978-3-540-79236-9_7
- Johnston EL, Dafforn KA, Clark GF, Rius M, Floerl O (2017) How anthropogenic activities affect the establishment and spread of non-indigenous species post-arrival. *Oceanography and Marine Biology Annual Review* 55: 389–420
- Jones PD (2000) The Mersey Estuary – Back from the Dead? Solving a 150-Year Old Problem. *Water and Environment Journal* 14: 124–130, <https://doi.org/10.1111/j.1747-6593.2000.tb00238.x>
- Keane RM, Crawley MJ (2002) Exotic plant invasions and the enemy release hypothesis. *Trends in Ecology & Evolution* 17: 164–170, [https://doi.org/10.1016/S0169-5347\(02\)02499-0](https://doi.org/10.1016/S0169-5347(02)02499-0)
- Keller RP, Drake JM, Drew MB, Lodge DM (2011) Linking environmental conditions and ship movements to estimate invasive species transport across the global shipping network. *Diversity and Distributions* 17: 93–102, <https://doi.org/10.1111/j.1472-4642.2010.00696.x>
- Kim TW, Micheli F (2013) Decreased solar radiation and increased temperature combine to facilitate fouling by marine non-indigenous species. *Biofouling* 29: 501–512, <https://doi.org/10.1080/08927014.2013.784964>
- Kolar CS, Lodge DM (2001) Progress in invasion biology: predicting invaders. *Trends in Ecology and Evolution* 16: 199–204, [https://doi.org/10.1016/S0169-5347\(01\)02101-2](https://doi.org/10.1016/S0169-5347(01)02101-2)
- Levine JM, Adler PB, Yelenik SG (2004) A meta-analysis of biotic resistance to exotic plant invasions. *Ecology Letters* 7: 975–989, <https://doi.org/10.1111/j.1461-0248.2004.00657.x>
- Lockwood JL, Cassey P, Blackburn T (2005) The role of propagule pressure in explaining species invasions. *Trends in Ecology & Evolution* 20: 223–228, <https://doi.org/10.1016/j.tree.2005.02.004>
- Macreadie PI, Fowler AM, Booth DJ (2011) Rigs-to-reefs: will the deep sea benefit from artificial habitat? *Frontiers in Ecology and the Environment* 9: 455–461, <https://doi.org/10.1890/100112>
- Madhani JT, Brown RJ (2015) The capture and retention evaluation of a stormwater gross pollutant trap design. *Ecological Engineering* 74: 56–59, <https://doi.org/10.1016/j.ecoeng.2014.09.074>
- Maguire RJ (2000) Review of the persistence, bioaccumulation and toxicity of tributyltin in aquatic environments in relation to Canada's Toxic substances Management Plan. *Water Quality Research Journal of Canada* 35: 633–679
- Mayer-Pinto M, Johnston EL, Bugnot AB, Glasby TM, Airoidi L, Mitchell A, Dafforn KA (2017) Building “blue”: An eco-engineering framework for foreshore developments. *Journal of Environmental Management* 189: 109–114, <https://doi.org/10.1016/j.jenvman.2016.12.039>
- Miller RJ, Etter RJ (2008) Shading facilitates sessile invertebrate dominance in the rocky subtidal Gulf of Maine. *Ecology* 89: 452–462, <https://doi.org/10.1890/06-1099.1>
- Minchin D, Gollasch S (2003) Fouling and ships' hulls: how changing circumstances and spawning events may result in the spread of exotic species. *Biofouling* 19: 111–122, <https://doi.org/10.1080/0892701021000057891>
- Minchin D, Nunn J (2014) The invasive brown alga *Undaria pinnatifida* (Harvey) Suringar, 1873 (Laminariales: Alariaceae), spreads northwards in Europe. *BiolInvasions Records* 3: 57–63, <https://doi.org/10.3391/bir.2014.3.2.01>
- Mineur F, Cook EJ, Minchin D, Bohn K, MacLeod A, Maggs C (2012) Changing coasts: Marine aliens and artificial structures. *Oceanography and Marine Biology: An Annual Review* 50: 189–234, <https://doi.org/10.1201/b12157-5>
- Morris RL, Deavin G, Hemelryk Donald S, Coleman RA (2016) Eco-engineering in urbanised coastal systems: consideration of social values. *Ecological Management & Restoration* 17: 33–39, <https://doi.org/10.1111/emr.12200>
- Moyle PB, Light T (1996) Fish invasions in California: do abiotic factors determine success? *Ecology* 77: 1666–1670, <https://doi.org/10.2307/2265770>
- Naylor LA, Coombes MA, Venn O, Roast SD, Thompson RC (2012) Facilitating ecological enhancement of coastal infrastructure: The role of policy, people and planning. *Environmental Science & Policy* 22: 36–46, <https://doi.org/10.1016/j.envsci.2012.05.002>
- Oberndorfer E, Lundholm J, Bass B, Coffman RR, Doshi H, Dunnett N, Gaffin S, Köhler M, Liu KKY, Rowe B (2007) Green roofs as urban ecosystems: ecological structures, functions, and services. *Bioscience* 57: 823–833, <https://doi.org/10.1641/B571005>
- Ostroumov S (2005) Some aspects of water filtering activity of filter-feeders. *Hydrobiologia* 542: 275, <https://doi.org/10.1007/s10750-004-1875-1>
- Perkol-Finkel S, Ferrario F, Nicotera V, Airoidi L (2012) Conservation challenges in urban seascapes: promoting the growth of threatened species on coastal infrastructures. *Journal of Applied Ecology* 49: 1457–1466, <https://doi.org/10.1111/j.1365-2664.2012.02204.x>
- Pimentel D, Lach L, Zuniga R, Morrison D (2000) Environmental and economic costs of nonindigenous species in the United States. *Bioscience* 50: 53–65, [https://doi.org/10.1641/0006-3568\(2000\)050\[0053:EAECON\]2.3.CO;2](https://doi.org/10.1641/0006-3568(2000)050[0053:EAECON]2.3.CO;2)
- Piola RF, Dafforn KA, Johnston EL (2009) The influence of antifouling practices on marine invasions: a mini-review. *Biofouling* 25: 633–644, <https://doi.org/10.1080/08927010903063065>

- Piola RF, Johnston EL (2008) Pollution reduces native diversity and increases invader dominance in marine hard-substrate communities. *Diversity and Distributions* 14: 329–342, <https://doi.org/10.1111/j.1472-4642.2007.00430.x>
- Prosser T, Morison PJ, Coleman RA (2015) Integrating stormwater management to restore a stream: perspectives from a waterway management authority. *Freshwater Science* 34: 1186–1194, <https://doi.org/10.1086/682566>
- Rivero NK, Dafforn KA, Coleman MA, Johnston EL (2013) Environmental and ecological changes associated with a marina. *Biofouling* 29: 803–815, <https://doi.org/10.1080/08927014.2013.805751>
- Ruiz GM, Fofonoff PW, Carlton JT, Wonham MJ, Hines AH (2000) Invasion of coastal marine communities in North America: apparent patterns, processes and biases. *Annual Review of Ecology and Systematics* 31: 481–531, <https://doi.org/10.1146/annurev.ecolsys.31.1.481>
- Schiff K, Diehl D, Valkirs A (2004) Copper emissions from antifouling paint on recreational vessels. *Marine Pollution Bulletin* 48: 371–377, <https://doi.org/10.1016/j.marpolbul.2003.08.016>
- Sella I, Perkol-Finkel S (2015) Blue is the new green—Ecological enhancement of concrete based coastal and marine infrastructure. *Ecological Engineering* 84: 260–272, <https://doi.org/10.1016/j.ecoleng.2015.09.016>
- Serrano E, Coma R, Ribes M, Weitzmann B, García M, Ballesteros E (2013) Rapid northward spread of a zooxanthellate coral enhanced by artificial structures and sea warming in the western Mediterranean. *PLoS ONE* 8: e52739, <https://doi.org/10.1371/journal.pone.0052739>
- Shea K, Chesson P (2002) Community ecology theory as a framework for biological invasions. *Trends in Ecology & Evolution* 17: 170–176, [https://doi.org/10.1016/S0169-5347\(02\)02495-3](https://doi.org/10.1016/S0169-5347(02)02495-3)
- Sim VXY, Dafforn KA, Simpson SL, Kelaher BP, Johnston EL (2015) Sediment contaminants and infauna associated with recreational boating structures in a multi-use marine park. *PLoS ONE* 10: e0130537, <https://doi.org/10.1371/journal.pone.0130537>
- Simkanin C, Davidson IC, Dower JF, Jamieson G, Therriault TW (2012) Anthropogenic structures and the infiltration of natural benthos by invasive ascidians. *Marine Ecology* 33: 499–511, <https://doi.org/10.1111/j.1439-0485.2012.00516.x>
- Simkanin C, Dower JF, Filip N, Jamieson G, Therriault TW (2013) Biotic resistance to the infiltration of natural benthic habitats: examining the role of predation in the distribution of the invasive ascidian *Botrylloides violaceus*. *Journal of Experimental Marine Biology and Ecology* 439: 76–83, <https://doi.org/10.1016/j.jembe.2012.10.004>
- Tan ELY, Mayer Pinto M, Johnston EL, Dafforn KA (2015) Differences in intertidal microbial assemblages on urban structures and natural rocky reef. *Frontiers in Microbiology* 6: 1276, <https://doi.org/10.3389/fmicb.2015.01276>
- Terlizzi A, Frascchetti S, Gianguzza P, Faimali M, Boero F (2001) Environmental impact of antifouling technologies: state of the art and perspectives. *Aquatic Conservation: Marine and Freshwater Ecosystems* 11: 311–317, <https://doi.org/10.1002/aqc.459>
- Tilman D (1997) Community invasibility, recruitment limitation, and grassland biodiversity. *Ecology* 78: 81–92, [https://doi.org/10.1890/0012-9658\(1997\)078\[0081:CIRLAG\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1997)078[0081:CIRLAG]2.0.CO;2)
- Tyrrrell MC, Byers JE (2007) Do artificial substrates favor nonindigenous fouling species over native species? *Journal of Experimental Marine Biology and Ecology* 342: 54–60, <https://doi.org/10.1016/j.jembe.2006.10.014>
- Vergés A, Steinberg PD, Hay ME, Poore AGB, Campbell AH, Ballesteros E, Heck KL, Booth DJ, Coleman MA, Feary DA, Figueira W, Langlois T, Marzinelli EM, Mizerek T, Mumby PJ, Nakamura Y, Roughan M, van Sebille E, Gupta AS, Smale DA, Tomas F, Wernberg T, Wilson SK (2014) The tropicalization of temperate marine ecosystems: climate-mediated changes in herbivory and community phase shifts. *Proceedings of the Royal Society of London B: Biological Sciences* 281: 20140846, <https://doi.org/10.1098/rspb.2014.0846>
- Vitousek PM, D'Antonio CM, Loope LL, Rejmanek M, Westbrooks R (1997) Introduced species: a significant component of human-caused global change. *New Zealand Journal of Ecology* 21: 1–16
- Warnken J, Dunn RJK, Teasdale PR (2004) Investigation of recreational boats as a source of copper at anchorage sites using time-integrated diffusive gradients in thin film and sediment measurements. *Marine Pollution Bulletin* 49: 833–843, <https://doi.org/10.1016/j.marpolbul.2004.06.012>
- Wasson K, Fenn K, Pearse JS (2005) Habitat differences in marine invasions of central California. *Biological Invasions* 7: 935–948, <https://doi.org/10.1007/s10530-004-2995-2>
- Weis JS, Weis P (1992) Construction materials in estuaries: reduction in the epibiotic community on chromated copper arsenate (CCA) treated wood. *Marine Ecology Progress Series* 83: 45–53, <https://doi.org/10.3354/meps083045>
- Weis P, Weis JS, Proctor T (1993) Copper, chromium, and arsenic in estuarine sediments adjacent to wood treated with Chromated-Copper-Arsenate (CCA). *Estuarine, Coastal and Shelf Science* 36: 71–79, <https://doi.org/10.1006/ecs.1993.1006>
- Wilcove DS, Rothstein D, Jason D, Phillips A, Losos E (1998) Quantifying threats to imperiled species in the United States. *Bioscience* 48: 607–615, <https://doi.org/10.2307/1313420>
- Willan RC, Russell BC, Murfet NB, Moore KL, McEnulty FR, Horner SK, Hewitt CL, Dally GM, Campbell ML, Bourke ST (2000) Outbreak of *Mytilopsis sallei* (Recluz, 1849) (Bivalvia: Dreissenidae) in Australia. *Molluscan Research* 20: 25–30, <https://doi.org/10.1080/13235818.2000.10673730>
- Williamson M (1996) Biological Invasions. Chapman & Hall, London, pp 1–244
- Williamson M, Fitter A (1996) The varying success of invaders. *Ecology* 77: 1661–1666, <https://doi.org/10.2307/2265769>
- Williamson MH, Brown KC, Holdgate MW, Kornberg H, Southwood R, Mollison D (1986) The analysis and modelling of British invasions. *Philosophical Transactions of the Royal Society of London* 314: 505–522, <https://doi.org/10.1098/rstb.1986.0070>
- Wilson JC, Elliott M (2009) The habitat-creation potential of offshore wind farms. *Wind Energy* 12: 203–212, <https://doi.org/10.1002/we.324>
- Wilson JC, Elliott M, Cutts ND, Mander L, Mendão V, Perez-Dominguez R, Phelps A (2010) Coastal and offshore wind energy generation: Is it environmentally benign? *Energies* 3: 1383–1422, <https://doi.org/10.3390/en3071383>